




Anna Maria Owen



XIII —

957. e. 37.

Perfect - 2 -



A
New Treatise
OF
Canary Birds.



No.

(J. E.)

Herriena de Chanteloup



A 17 Va
New TREATISE
OF
Canary-Birds.

CONTAINING

The manner of Breeding and
Coupling them, that they may
have Beautiful Young Ones.

WITH

Curious Remarks relating to the
Signs and Causes of their Distem-
pers, and the Method of Curing them.

Written in *French* by Mr. *Hervieux*,
and Translated into *English*.

LONDON:

Printed for Bernard Lintot, at the
Cross-Keys between the Temple-Gates:
And Benjamin Barker, and Charles King
in Westminster-Hall, 1718.

A

NEW TREATISE

CANARY-BIRDS.

Jos. Banks

WITH

Several Remarks relating to the
Sings and Customs of these Birds;
and the Method of tending them.

Written in French by Mr. Huet
and Translated into English.

LONDON:

Printed for Bernard Lintot, at the
Grave-Knave between the Temple-Gate
and Bream's Church, and for the King
in the Strand, at the Sign of the Crown.



THE

Authors Preface:

Shewing how Necessary the Contents of this Treatise are, for such as desire to enjoy the Innocent Recreation of *Canary-Birds*.

IT is requisite, that those who are inclin'd to Enjoy the *Diversion of Canary Birds*, should be acquainted with their *Inclination* which differ very much among themselves; what Food is proper for them, and their *Dis-temperers*.

The Preface




stempers. By knowing their Inclinations, many Accidents that befall them, are prevented; especially when it is design'd to make them Sit. By knowing what proper Food to give them, according to the several Seasons, the Birds are always kept in perfect Health; and lastly, by understanding their Distempers, they are often rescu'd from speedy and inevitable Death. This is what I propose in the present Treatise of Canary-Birds. I am not ignorant, that a Treatise has been writ, of all Birds in general; but the Author of it has allow'd but one Chapter for the Canary-Birds; leaving it perhaps to some other, who is better acquainted to say more of them. I shall therefore add, at least 24 Chapters, to that which has been already

The Preface


already writ on this Subject. For observing the better Method, I cannot forbear repeating something that has been said already, but it would be needless for me to set Pen to Paper, if I had not something New to add to what has been made publick; and did not intend to Explain and Enlarge what has been written concerning the Canary-Birds. This Treatise will therefore be an Improvement of all that has been said on this Subject, and an Exposition of what has not been hitherto made very plain, being myself perfectly knowing in what I am going to Write about the Canary-Birds. I am satisfy'd others will take great Satisfaction in the Diversion these lovely Little Birds afford, they will not be surpriz'd at the several

The Preface



ral Accidents that may befall them; nor will they wonder to see them Dye, after having done all that they thought was requisite for prolonging of their Lives, knowing that all Living Creatures must sooner or later, have an End. I will endeavour to explain my self upon every particular, as intelligibly, and in as few Words, as may be.




B  L



Prelude



Marche des Surlobes



Handwritten musical score on six staves. The notation is in a historical style, featuring square notes and various clefs. The text "36. Cantata" is written at the top right. The text "Missa de S. Jo. Baptista" is written across the middle staves. The text "B L" is written at the bottom left, with a crown symbol between the letters.

36. Cantata

Missa de S. Jo. Baptista

B L



A
New Treatise
OF
Canary Birds.

C H A P. I.

Of the Original of the Canary Bird.

I Will not here spend my Time in repeating what several ancient Poets have said concerning the Original of the *Canary* Bird; because that Account is Fabulous. It shall suffice to say, that the first we ever saw in these parts, were brought us from the *Canary* Islands, sub-
B ject

ject to the Crown of *Spain*, and from thence they had the Name they are known by; tho' they are now also brought to us from *Inspruch*, the Capital of the County of *Tirol* in *Germany*, and that by *Germans* twice a Year; that is, in the Spring, and in Autumn, of which more shall be said hereafter. For my own part, I now call them *French*, or *English*, and not *Canary* Birds, because having been here so many Descents they must be look'd upon as Naturaliz'd. I shall add no more concerning their Original, leaving that to some other; who when he has much search'd in vain after these little Creatures, will find no Man to justify what he may advance concerning them.

CHAP. II.

The Names given to Canary-Birds, according to their several Colours.

I Think it will be proper in the next place, to set down the Names commonly given to *Canary-Birds*, according to their several Colours; to the end that every

of Canary-Birds. 3

every one may know, in what Class, or rather of what degree of Beauty the Birds are he has, or designs to Buy; and therefore I shall name them in their Order, beginning with the most Common Sorts, and concluding with the most Beautiful.

Common Grey *Canary*-Birds.

Grey *Canary*-Birds, rough footed and the Feet white, call'd of the Coppel-crown breed.

Grey *Canary*-Birds with white Tails, of the Coppel-crown breed.

Common Ash-colour *Canary*-Birds.

Ash-colour *Canary*-Birds with red Eyes.

Ash-colour *Canary*-Birds, with a cast of Gold colour.

Rough-footed Ash-colour *Canary*-Birds, of the Coppel-crown breed.

Ash-colour *Canary*-Birds, with white Tails, of the Coppel-crown breed.

Common Lemon-colour *Canary*-Birds.

Rough-footed Lemon-colour *Canary*-Birds, of the Coppel-crown breed.

Lemon-colour *Canary*-Birds with white Tails, of the Coppel-crown breed.

Common Mottle *Canary*-Birds.

Mottle *Canary*-Birds with red Eyes.

Mottle *Canary*-Birds with white Tails, of the Coppel-crown breed.

Common Buff-colour *Canary*-Birds.

Buff-colour *Canary*-Birds with red Eyes.

Buff-colour *Canary*-Birds, with a cast of Gold-colour.

Buff-colour *Canary*-Birds, rough footed, of the Coppel-crown breed.

Buff-colour *Canary*-Birds, with white Tails, of the Coppel-crown breed.

White *Canary*-Birds with red Eyes.

Common Coppel-Crown *Canary*-Birds.

Coppel-crown *Canary*-Birds with red Eyes.

Canary-Birds with Ash-colour Coppel-crowns.

Canary-Birds with Ash-colour Coppel-crowns and red Eyes.

Canary-Birds with Black Coppel-crowns.

Canary-Birds with Black and Lemon-colour Coppel-crowns and red Eyes.

Canary-Birds with regular Black and Lemon-colour Coppel-crowns.

These are the common Names given to *Canary*-Birds, they being generally call'd and distinguish'd by their Colours.

It remains that I mention the Names of those which are call'd Mongrel *Canary*-Birds; which are the Breed of Cock or Hen *Canary*-Birds, which have been coupled with Birds of a different Kind, as Yellow-Hammers, Chaffinches, Linnets, Goldfinches, or the like; then those that come of them take their Names from the sort of Bird they were coupled with; as
for

of Canary-Birds. 5

for example, when a Cock *Canary-Bird* is coupled with a Hen-*Chaffinch*, the Young Ones which come of them are call'd *Mongrel Canary-Birds* of a *Chaffinch*, and so the others.

Mongrel Canary-Birds of a *Linnett*.

Mongrel Canary-Birds of a *Yellow-Hammer*.

Mongrel Canary Birds of a *Gold-finch*, and so of others.

CHAP. III.

Of the Cages and Breeding Places fit for Canary-Birds.

THERE are abundance of different Sorts of Breeding-Places continually made, according to every Mans Fancy and Ability. Those who make them of Deal have them very Cheap; but it is certain, that when the Birds have Hatch'd in them one Year at farthest, they are fit for nothing but to Burn, as well on account of the many small Insects or Worms they breed, as because of the slightness of that Wood, which cannot be long expos'd to the Air, but it will warp and

yield every way : Those which are made of Beech are somewhat better, that Wood being closer and more solid than Deal ; and tho' it is of no better Colour it is Dearer. But those who would have neat and more lasting Huts or Breeding-places, must make them of Wainscott, which is a stronger sort of Wood, and of an agreeable Colour ; and the older it is, the harder and more beautiful it grows.

Lastly, Those who will have these Huts in Perfection, must cause them to be made of Walnut-Tree, a Beautiful and Serviceable sort of Wood ; and the Bottoms and Drawers must be all of a piece, that they may not give way by Length of Time. They may depend upon it, that the longer they keep them, the handsomer they will look ; and that if they Live never so long they will not see the end of them. For my own part, my Method is to have these Huts quite open ; that is, all the four Sides of them to be of Brass Wire, and that for two principal Reasons.

1. Howsoever these Huts are plac'd in a Room, the Birds are always in Sight, which way soever a Man looks at them, which is very agreeable and pleasing to the Eye.

2. Because those Birds, who are so expos'd to view, become so Tame by seeing People

People on every side, and at all times, that nothing scares them, and consequently they are not apt to beat themselves by fluttering, when they are Dress'd, as often *Canary-Birds* do that are kept in Dark Cages. I will not here speak of those costly and rich Huts that may be made, as some Persons in this City have them, the Bottoms and two Sides whereof are Looking-Glass, and the Borders and Corners Painted with several sorts of Birds in Miniature, to the end that the *Canary-Birds* kept in those Fine Huts, may have their Young, if possible, like those there Painted. I leave it to others to make Reflections on those extravagant Expences, which might serve to Relieve many Families, whom the Difficulties of these Times have reduc'd to extream Want; I shall only say that it were practicable to imitate those Fine Huts, without being at the same Expence, by placing large plain Glasses, instead of those Looking Glasses; I have spoken of, and caus'd some Birds of several Colours to be Painted on those same Glasses, to cause an Impression on those *Canary-Birds*, that they may breed such Young as are there Painted.

As for Cages, I will add a word concerning them by the way: The round are very prejudicial to the Birds, because they have not room enough to walk, and so they are

always har'd. The long Cages are best, not quite so broad, but lofty. A Bird in such a Cage is not har'd, having room to fly in the Height, and to walk in the Length, and therefore grows stronger and healthier than others; and being easy in his Confinement is never subject to Melancholy, which proves a fatal Distemper to them. I have contriv'd a sort of Cage, which pleases me very well, and has been approv'd of by Judicious People; it is long and proportionably broad, and of a good Height; there are no Troughs on the sides; so that the Bird is plainly to be seen at any distance from the Cage; whereas the two Troughs commonly plac'd on the sides of common Cages, do so cover the Bird, that when he is eating, or stands on the Perch next to those Troughs, he is not seen, especially when we are at a distance from the Cage, or sitting.

The two Leaden Pots in this new sort of Cage, are plac'd underneath, fix'd into the Drawer at the end of the Cage; so that in taking out of the Board, which is done at the back part of the Cage, you at the same time take out the two Troughs in which the Birds Meat is. The Troughs are wir'd before, at proper distances, within the Cage, to the end that the Bird only putting in his Head, may

Canary-Birds.

9

may not throw out the Seeds ; which is not so in the other Cages.

The advantage of these new Cages is, in the first Place, the Pleasure of seeing your whole Bird at any Distance, there being nothing to obscure him which way soever he moves in the Cage.

Besides, the Bird that does not continually see his Meat, when he is upon his Perches, does not eat so often, and therefore does not grow so Fat, but more beautiful, and sings oftner, and is not so apt to gorge himself, which they are subject to, by too much eating, and scarce ever escape alive, when it happens.

This sort of Cage is also very advantageous to them when they are Sick, or their Feet are hurt, for then they can come at their Meat as they stand down, without getting upon their Perches, which is not so in the other Cages, where they are often found Dead under their Troughs, for want of Strength to get upon their Perches to feed. There are many more Conveniences belong to these Cages, which would take me up too much time to enumerate.

For the better conceiving of this sort of Cage, I have thought fit to give a Draught of it.

A is the Front of the Cage, B the Back Part of it, C the Trough, D the Top of

B 5

the

the Trough, E a little Board which is the Front of the Trough, made fast to the lower edge of the Cage, to keep the Trough steady, and support the Wires made fast to the top of the Trough, F the Board to draw out of the Cage, with the Troughs on it.

I will not spend time in mentioning the several ways of adorning Cages, Voleries and Huts, daily invented by several curious Persons for their *Canary*-Birds. Some make the Frames of their Cages or Huts of Box, and instead of Brass use Silver or Gold Wire; others adorn them with false Diamonds; some embellish them with Amber; others inlay them with Ivory, all according to an exact Symmetry. In short, every nice Person does it according to his Purse or his Fancy.

CHAP.

CHAP. IV.

*The proper Season for coupling of
Canary-Birds, when it is de-
sign'd to have a Breed, and
of the most advantageous Posi-
tion.*

AS to the time of coupling them, in order that they should multiply, that cannot be exactly determin'd; but the Season is to be observ'd, which some Years is sooner or later than others. When you perceive, that the Sun begins to grow warm and the nipping Frosts are over, which is commonly about the latter end of *March*, you may prepare to couple your *Canary-Birds* after this manner.

You must take a new Cage, or at least one that is very clean, that there may be no Worms in it; into which put a Cock *Canary-Bird* with the Hen you have design'd for him. They grow acquainted and couple sooner in a little Cage than in a great Hut, as being closer confin'd and nearer

nearer one another. Special Care must be taken, not to do, as happens every Year to many, who put two Cocks or two Hens together, for not having parted the Males from the Females in due time; thus they are often confounded when the coupling time comes; for some Hens in the Spring sing almost as loud as the Cocks; and on the contrary, some Cocks sing so low, that they are often mistaken for Hens. When a Man has committed this mistake it makes him fret, for as it often happens that one of the two Hens, which have been put together lays, it is evident that the Eggs cannot hatch, and then the Fault is laid upon the Cock, which in reality is no other than a Hen. On the other Hand, if you put two Cocks together, for want of knowing them, then are you no less concern'd to see that the Bird you take for a Hen does not lay; then the pretended Hen is accounted Barren, and so the Year passes away with this Disappointment, without discovering the Mistake; and the reason why it is certainly look'd upon to be a Hen, is because that Cock which is taken for a Hen, sings very little or not at all, and that nothing extraordinary; for when you put two Cocks into a Cage, it frequently happens that one of them either through Fear, or otherwise, does not sing. Therefore, when you have kept your Pair
of

of *Canary-Birds* 8 or 10 Days in a little Cage, and you perceive that they are well match'd, as plainly appears, when they do not Fight, as they commonly do the first Days they are put together, and that on the contrary they make Love and Bill, then you may let them into the Hut design'd for them, which is to be made as describ'd in the 3^d Chapter, and you must there give them all that is requisite for making of their Nests, of which I shall treat hereafter.

As for the Position of the Place, I must own they sit wheresoever they are plac'd, either facing to the East or West, the North or South, in the Town or Country, within Rooms or without, that is, on Windows that are open all Night; but I deny that the young ones will thrive as well in one Place as in another. If therefore you would have thriving *Canary-Birds*, you must place their Huts fronting to the East, above any other Quarter. There the old ones will be less liable to several Accidents and Diseases which befall them, when they are ill posited; the new hatch'd Birds thrive more in one Day, when their Position is advantageous than they would otherwise do in two; for if they are exposed to the West or South, it burns their Brains, and breeds abundance of Worms, and often causes the Hens to sweat, and so they

they stifle the young. Sometimes there comes such a cold Wind from the North, even in Summer, as kills the new hatch'd Birds, and sometimes the old ones. I will not discourse of many other unlucky Accidents that befall them, as either not laying at all the whole Year, or else laying only of addle Eggs, sometimes for being posited in such manner as does not suit with them, or else for being in a Place too dark, which makes them Melancholy, and breeds Sores. It would be endless to mention all the ill Accidents that befall *Canary-Birds* in their Huts, which commonly proceed from having plac'd them to sit in an Air, which is not agreeable to their tender Disposition. I believe those who are acquainted with *Canary-Birds* will approve of what I have said to this purpose.

CHAP. V.

The certain Method for coupling of Canary-Birds that the young may be of beautiful Colours.

THE more *Canary-Birds* have multiply'd in this City, and are consequently grown the more common, the more nice People are grown, as to their Colours; those who ten Years ago were well pleas'd with a Grey *Canary-Bird* of two Pistoles Price, will not be now satisfy'd with the common cottle Crowns; the common Ash and Gold Colour, the Buff and Mottled are now of little Value among the curious; they will have their *Canary-Birds* as pleasing to the Eye by their Plumage, as they are to the Ear, for their sweet warbling. I have therefore here set down how *Canary-Birds* are to be coupled to expect the young ones more beautiful than the old. I begin with the most common Breeds, and will conclude with the beautifullest we have at this Time.

They

They who couple a Grey Cock and a Grey Hen, which are both common, can expect none but a Grey Breed.

The same is to be said of the Ash, Buff, Mottled and Lemon, coupled with Hens of the same Colour, which cannot produce any Birds more beautiful than themselves.

But when the different Kinds are mix'd, it falls out better, for Nature often delights in producing finer and more beautiful Birds than was expected.

It is not always requisite to have copple Crown *Canary*-Birds in order to make a beautiful Breed; but it is enough that they be of the copple Crown Breed, for those that come from them are often more beautiful than if they came directly from Copple Crowns.

For Instance, A white Tail Grey Cock, with a rough footed Grey Hen, may produce, besides the rough footed Greys with white Tails, which are to be expected of course, some with Copple Crowns, often more regular than if they had been bred from Copple Crowns. The same may be said of an Ash, Lemmon, Buff and Mottled Cock, which being of the Copple Crown Breed, and that is known by their being rough footed, or having some white Feathers in their Tails, being coupled with Hens of their own kind, Breed beautiful

tiful Birds and often Copple Crown'd.

But those who would have still more beautiful *Canary-Birds* must Match them thus.

A Cock with an Ash Colour Copple Crown, and a Lemmon Hen, with a white Tail, will make a beautiful Breed.

All Copple Crown Cocks, with white Tail Hens, except the Grey with white Tails, produce very fine Birds.

They who join Copple Crown Cocks and Hens, will have all Copple Crown young, tho' sometimes they happen to be Grey; and that is because either the Dam or the Sire of that pair of Copple Crown *Canary-Birds* was Grey. But to make short, they that would have some of that fine Breed call'd the Jonquille with a black Copple Crown, which is at this Time reckoned the most beautiful and highest valu'd sort, must put together a Jonquille, or Lemmon Colour *Canary-Bird* of the Copple Crown Race, and a Jonquille, or Lemmon Hen.

If you would have more of the yellow and more of the Copple Crown, you must on the contrary couple a Cock with a black Copple Crown, and a yellow Hen with a white Tail, which produces the beautiful Jonquille; but that this may succeed the better, the yellow Hen with a white Tail, here spoken off, must be from a Jonquille
Cock

Cock well mark'd, and from a yellow Hen with a white Tail. In a Word, this is all that can be done towards producing Birds of beautiful Colours. The young which come from this last Race I have here spoken of, are much more troublesom to rear than all other sorts, as being of a very tender Constitution.

CHAP. VI.

What Things are necessary for making the Canary-Birds Nests.

THERE are Seven or Eight several sorts of Things commonly given to *Canary-Birds* to make their Nests, as new or common Deers Down or Wool, Hay, Moss, Cotton cut small, coarse Hemp, or Flax, Soft long Hair of Fur, &c.

Of all these several things, only one or two ought to be us'd for making of the *Canary-Birds Nests*, all the rest being very improper, as I shall next observe.

For instance, Cotton cut small, as well as Flax, is apt to cling to their Claws, whence it often happens, that the sitting Hen, rising hastily from off her Eggs, drags

drags the Nest after her, and consequently breaks the Eggs that are in it, and this may be done without perceiving the Cause of it; then a Man is apt to look upon it as the Fault of the Cock or the Hen, which are suspected to have done it designedly, and yet you see they knew nothing of it. New or common Deers Down or Wool is not so good for them as is generally imagin'd, for it puts the fitting Hens into such a Heat, that they often sweat, and when the young ones are hatch'd, they are stifled by it in a few Days. Besides, that Wool being heated, sticks so close to the Bodies of the new hatch'd Birds, like a Paste, that they can not void their Excrement, and thus dye, with their Crows full, so that the Cause of their Death is not perceiv'd. If I were to make use of that Down, it should be only of the new, and would only employ it about their first Nest, because it is not yet hot Weather, when they have their first Breed, but it is never to be given them at their second, third and fourth laying, to avoid the ill Accidents I have mention'd. They must also be allow'd no Moss, or very little; for when they have much, it sometimes happens that they bury their Eggs, so that they lying at the Bottom of the Nest and the Moss over them, a Man is apt to believe the
Hen

Hen has not laid, when she has several Eggs.

They must therefore have nothing given them but very fine and small Hay, to make the Body of their Nest; besides that Hay must be gather'd and well dry'd in the Sun some time before they have it, that the strength of it being gone, it may not affect their Heads. When the Nest is almost made, there may be given them a little Pinch of Moss, well dry'd in the Sun, and the same Quantity of Deers Wool, which must be allow'd only for the first Nest and not for the others, for the Reasons above mention'd.

There is a sort of fine furry Hair, which may be had at the Brush-makers, and is very proper for these Birds. You may take the finest and shake it very well, to beat out the Dust, or if you would do better, wash and then dry it in the Sun, which will take out all the Dust and the Scent it naturally has, then strew it about the Cage; you will be pleas'd to see what a charming Nest they will make. This Hair alone, without any other thing will serve them to make their Nest, and that which has already serv'd in one Nest, being again wash'd in boiling Water, will be fit for another Nest.

There are three sorts of things given them to make their Nests in; first little
Oser

Canary-Birds.

21

Offer Baskets. 2. Wooden Cases, or Boxes. 3. the like made of Earthen Ware.

Those who invented the Earthen Pots, say they did it, because the Hen *Canary*-Bird, who is at least 24 Days without coming out of her Nest, does not sweat in them as she often does in the Baskets; but in my Opinion they have not succeeded in that Point; for it is certain, that the Earthen Pot, being naturally damp, must have ill effects upon the Sitting Hen, and worse upon the little new hatch'd Birds.

One of the most considerable is, that if the Hut happens to be a little expos'd to the Sun, the Earthen Pot will grow so hot and retain the Heat so long, even after the Sun is gone off it, that it will certainly kill the Hen and stifle the young ones.

As for the Wooden Boxes or Cases, they are convenient, because it is easy in them to take out the Nests and put them in again now and then, to shake out the Insects that lye at the bottom; for in that sort of Boxes the Nest is taken out altogether, and may be put in again exactly as it was before, without the Birds perceiving that it has been touch'd; but there is no making use of those Boxes that are bor'd through, because in taking out the
Elder

Elder Stick that goes across them, the Nests are all disorder'd and the Eggs often broken; which the Hens perceiving, they generally take Pet, and leave their Nests.

Two other Inconveniencies attend the wooden Boxes. 1. The Nest which is made in a wooden Box having no passage for the Air, grows very hot, and often causes the Hen to sweat. 2. The Nest made in those Boxes is so small and close, that when either the Cock or the Hen gets into it they drag it with their Claws and break the Eggs, or overturn the little ones, if there are any in it.

It is needless to say any more to this effect; but it is best, according to ancient Custom, to make use of little Wicker Baskets, in which the Nest is not so much stiv'd up and holds faster than in any other thing. Those who make the Baskets bigger than was formerly us'd are in the wrong, for the Birds spend more time in filling them, to make their Nests, and consequently toil themselves the more. Besides, in a great Basket, the Eggs a Hen sits on are apt to lye at a distance from one another, and it often happens that several of those Eggs do not hatch; because they were not well cover'd.

When they are in the Hut great care must be taken to give them very fine

River

River Sand, thorough dry, and therefore I would advise to sift it, to the end that when a Hen has a Mind to lay on that Sand, as it often happens, the Egg may not be broken. It also happens, that the Cocks and the Hens coming out of the Nest drag out after them the new hatch'd young ones, as I observ'd before, and they falling on a very fine Sand and clear from Stones, are not kill'd, as I have found by my own Experience.

Note, that they must have but one Basket given them at a time to make their Nest; for as has been well observ'd, when they have two given them they sometimes carry into one and sometimes into the other and so they do nothing to the purpose; whereas, when they have but one given them, they make their Nests orderly at once, and think of nothing but hatching their young. Twelve Days after the young are hatch'd, you must put in another Basket on the other Side, for they never fail to make their second Nest, tho' they are at the same time breeding their young. For my part I make their Nest my self, especially the Second, Third and Fourth, and only allow them to make the First. That fatigues them the less, especially the Copple Crowns; and if it happens not to please them, they have less trouble to mend it, than they would have

have to make the whole. My *Canary-Birds* like it very well, for they love to have their Work done for them.

CHAP. VII.

The several sorts of Food to be given to Canary-Birds, when they are pair'd, when they have young, and when they are in Cages.

THAT which commonly discourages most of those who first undertake to breed *Canary-Birds*, is the great Number of them that dye upon their Hands; which commonly happens through over or under-feeding them, without considering, that what is proper for them at one time, is often destructive at another. When therefore your Birds are well rear'd by the Hen, which you will know, by their feeding alone, you must give them for their common Food, Rapeseed, Miller, Canary-feed, and Hempseed, which are to be thus mix'd, *viz.* half a Pint of Hempseed, the same Quantity of Canary-feed, a Pint

a Pint of Millet, mix all with fix Pints of Rapeseed, well cleans'd, that there may be no Dust in it. Put this mixture into a close Deal Box, that no Dirt may get into it: Then fill the Birds Trough with this Grain, enough in Quantity to serve them two Days at least, to the end that when they have eaten the white the first Day, they may feed on the black the second, which will prevent their growing too Fat, and they will Sing the better, and this mixture of Seeds you shall continue as long as they live.

Some give them nothing but Rapeseed. Those Birds which can accustom themselves to this hard Diet, they say, live longer than others, but it occasions many of them to dye; for I have observ'd, that most of them grow so Lean and Melancholy, especially those that come of the last Sitting, which are generally weaker than the others, that the first Distemper carries them off.

Another Inconveniency of this hard feeding is, that when you put this sort of *Canary-Birds* into Huts, when it is requisite to give them other more substantial Food, especially when they have young ones, they eat so much of that new sort of Food, being before kept Starving, that they Surfeit and Dye in a few days; great care is therefore to be taken what Food

the *Canary-Birds* you Buy have been rear'd with and are actually fed with at that time; because several curious Persons, have different ways of feeding them; some, as has been said, giving them nothing but Rapeseed alone, and others allowing them much Millet, Canary-feed and Hempseed, with their Rapeseed, and then being ignorant of the common Food of those Birds you Buy, you are apt to give them that which is not proper; and that Change of Food making a strange Alteration in their little Bodies, ruins their Constitution, and impairs their Health, so that they often fall Sick and Dye, without your being able to guess at the cause.

It is therefore safer to have Birds at the first Hand, that is, of Townsmen, who have a mind to part with them; for they freely tell you what Food they have made use of in rearing the Birds you have bought, and by continuing the same Food you are out of Danger of losing them; whereas those who make it their Business to Trade in them, not being able to tell you what Food those you Buy have been rear'd with, as not knowing it themselves, being almost ignorant whence they had them, because they gather them from several Places, and do not take the trouble to enquire after it, as being likely to Sell them in a few days, and therefore during the short
time

time they have them in keeping, they spoil them themselves, by giving them three times more Food, than they have occasion for; and hence it follows that their Birds are carry'd to them Dead three Days after they were Sold, and when the Buyer complains, their Answer in short is, *that all Creatures must Dye as well as Men.*

When they are pair'd and put into Huts, you must sometimes give them, besides their ordinary Seeds I have here mention'd, some little Dainty, as for instance, a bit of Cake or hard Bisket, especially when you find the Hen is ready to Lay. You must also give them, during the first eight Days of their being in the Hut, much Lettice-seed, which purges and makes them evacuate the ill Humours they have contracted during the Winter.

We are now come to the most intricate and difficult time of managing *Canary-Birds*, which is, when they have young ones. Every Man then uses them according to his Fancy and therefore many do not prove successful. It is thus I use them at that time.

The Day before the young ones are to be Hatch'd, which is the 13th Day of the Hens sitting, I change their Sand, putting in fresh, clean all their Perches, fill their Trough with Seed, raking away what was

in it before, I also give them fresh Water, after making their Pot very clean, and all this to avoid disturbing of them during the first Days after their Young Ones are hatch'd. I give them also half a Simnel, taking off the upper Crust, and a little Bisket, both very hard, because if either of them were soft, they would eat too much, and then drinking upon it, would certainly be choak'd. As long as that Simnel and Bisket last, no other must be given them; but that which follows must be chang'd twice or thrice a Day, especially in very hot Weather, *viz.*

A Quarter of an hard Egg, cut very small, the White and the Yolk together; a bit of Simnel dipp'd in Water and squeez'd in the Hand, all of it laid on a little Saucer, and on another they must have of their common Seed, that has been steep'd two Hours, some Water must be thrown on it, and it will be better if the Seed be just boil'd up, and afterwards wash'd in cold Water, which takes off the Strength and Sharpness of the Seed. They eat it well, and do not gorge themselves, nor are the Young Ones in danger of being choak'd, tho' the Old ones give it them never so often.

They must also have some Greens given them, but a small quantity; *viz.* Chickweed, Groundsel, and when there is none left

left on the Ground, as in *July* and *August*, instead of them, the Heart of a Cabbage Lettice, a little Endive and a little very ripe Plantain. This must be fresh three times a Day; that is, once at 5 or 6 in the Morning, again at Noon, and lastly at 5 in the Evening. The Old must be taken away every time fresh is put in, because it all spoils in a few Hours, especially during the great Heats. For Instance, the Rapeseed grows sour and shoots, the wet Simnel grows sour, the Chickweed, Groundsel and Lettice, wither; and the Old ones giving the Young that decay'd Food, hinders them from thriving, and very often they Dye for want of Care, and of good fresh Food.

I give them besides, now and then some Lettice-seed, Pink Seed and Tansy-seed, all mix'd in a Pot, and I particularly observe, what the Cock likes best of all I give him, and when I have found it, I allow him as much as he will have, for when they have Young Ones, there is no reason to deny them what they love best, except the Greens, of which I do not allow them their fill; because as long as they have them they eat them rather than any other thing, and feeding their Young with nothing but those Greens, they give them an ill Habit of Body, and they often dye of that Excess. I also sometimes

put a bit of fresh Liquorice into their Water, which gives it a relish and does not heat them as Sugar does. In short, by this Method, which seems to be easy enough, I make many envy me, as saving more *Canary-Birds* than many do, who pretend to have been several Years well skill'd in this Art. I know some who give their *Canary-Birds* fifty other sorts of things, and yet they Dye sooner, and in greater Numbers, than those which are serv'd as I have here set down; and therefore I will not speak of those several sorts of Food, which are more hurtful to them, than proper to preserve their Health. In the hot Weather you must not forget to put into the Hut some fresh Water in a little Cup or Vessel, for the *Canary-Birds* to wash themselves, and the same must be done with the Young Ones, when they come to feed themselves. Take heed not to give them a Vessel too deep to wash in, lest they be in danger of Drowning. Change that Water every day in very hot Weather, as well as that in their Drinking Glass, or Pot.

CHAP. VIII.

Several sorts of Paste for Young Canary-Birds, when they are to be fed by Hand, or with a little Stick.

Several Persons make divers sorts of Paste to feed the Young *Canary-Birds*, when they are to be Fed by Hand, or with a little Stick; but some burn their Bowels by making the Composition too nourishing; and others, on the contrary by giving them a sort of common and too liquid Sustenance, weaken their *Canary-Birds* so much that there is no saving them from Death.

When therefore you design to feed your *Canary-Birds* by Hand, you must first observe, whether they are strong enough to be taken from under the Hen; for if you take them away too soon, they commonly decline every Day; and notwithstanding all your Care they fall away and Dye. Neither must they be left too long under the Hen; for when they are grown too

C. 4. big

big under her, they begin to know their Sire and Dam, grow fullen, and will not Feed; and though they be cover'd and put into a Dark Place, to make them forget the Cock and Hen, all these Precautions often prove useles, they pine and at last they must be return'd to the Cock and Hen to save them. The *Canary-Birds* you would feed by Hand must be well fledg'd. If they are Grey, or Ash, they may be taken from under the Hen, at 10, or 12 Days old, as being of a stronger Breed; if they are Copple Crown'd, they must not be taken away till 13 Days old; and lastly, if they are Lemmon colour, they must not be remov'd till 14 Days be past. This Rule I prescribe is not always without Exception, because sometimes there happen unlucky Accidents, which make it necessary not to follow it.

For instance, a Hen falls sick the 5th or 6th Day after the Young are hatch'd, in which case, there is no doubt but it is safer to take away the Young Ones to feed them with a Stick; but this is in case you have not another Hen, under which they may be put; which is better than to leave them to that Hen, which communicates her Distemper to those new hatch'd Birds, and Starves them for want of sufficient Food.

It also sometimes happens, that the
Hen

Hen feeds her Young so ill, that they visibly pine away, in which Case they must be sooner taken away than if they were well fed. Sometimes the Huts are so dark, and so ill posited, that the Little Ones do not thrive, tho' the Hen feeds them sufficiently. I have seen *Canary-Birds* a Fortnight old, which no Man would have judg'd to have been hatch'd above a Week; and the Fault was only to be ascrib'd to the Position of the Place, which was dark, close, and where the Sun, which chears their Bodies, never came. In this case they may also be taken away sooner than the common Rule prescribes, in order to feed them by Hand.

It often happens that the Hen forsakes her Young after 7 or 8 Days; then the Cock takes Care to feed them; but whatsoever different things can be given the Hen, to make her Second Nest, she will then pull the little Feathers off those Young, which kills them in a few days; and this is also a sufficient reason not to leave them any longer with the Old ones. There are many other reasons, too long to be here mention'd, which oblige to take the Little ones from under the Hen, to breed them up by Hand; but I advise to leave them under the Cock and Hen as long as I have prescrib'd, when there is no urgent Occasion to remove them; be-
C 5. cause.

cause the Food the Cock and Hen give them, strengthens and agrees with them better, than what a Man can prepare for them, be it never so good; and when they are taken away somewhat stronger, they are in a better Condition to withstand the Change of Diet, which is made when they are bred by Hand.

Here follow two sorts of Compositions that may be given them.

A sort of Paste for Canary-Birds, which may serve a Fortnight at least, without being spoil'd.

Take half a Pint of Rapeseed and bruise it very well in a large Mortar, or on a smooth Table, running a wooden Rouler over it several times; so that the Seed being well broken, you may cleanse it very well from the Husks; add to it three little dry Simnells, bruis'd and reduc'd to Pouder, after having taken off the first Crusty part; add to it a Penny Bisket; all this being well mix'd together and reduc'd to Pouder, put it into a new Wainscot Box, and set it in a Place that is not expos'd to the Sun. Take a spoonful, or more of this Pouder, as you have occasion, and your Birds Food will be made in a Moment, adding to it a little of the Yolk of an Egg and a Drop of Water, to moisten the whole.

But you must not give your *Canary-Birds*

Birds this Pouder above twenty Days at most; for I have observ'd, that after the said Term, the pounded Rape-Seed which is in the Composition, tho' dry, grows sour, and when Water is put to it has a Taste like Mustard, which cannot but be hurtful to your Young Birds. You are not therefore to follow the Notion of that Person, who has writ that this Composition would keep a long time, tho' kept in a Deal Box, a sort of Wood, which by reason of its Softness, is apt to breed Worms very soon, especially when any Pouder I have spoken of is in it. After 20 Days at most, if you have any of this Composition left, you may give it dry to the Old Cocks and Hens, who will eat it heartily, without receiving any Damage by it, and make fresh for those you bring up by Hand.

For my own part I own, I choose rather to be at the Trouble of making it fresh every Day, and am, perhaps with good reason, of Opinion, that the Young Birds, which have fresh Food made every day thrive better.

I make my Composition after this manner.

The three first days that I begin to Feed my Young Birds, I take a bit of Simnel, the crust whereof is taken off, because it has a sort of Bitterness; I add to it a very

ry little bit of Bisket, all hard, and reduce them both to Pouder; then I put to it half the Yolk of a hard Egg, or more if need be, which I moisten with a little Water, tempering all well, that there may be no hard Lumps. This Paste must never be too thin; because, if so, it does not nourish them so well, and they are continually craving; besides, that when it is too wett they are often Loose, and scarce recover it; But when the Paste is somewhat stiff, it lyes longer in their Crows, and nourishes them better: when the hard Egg is new, the White dissolves as well as the Yolk, and does not heat them so much, as if there were only the Yolk.

When the first three Days are past, and I perceive my Young Birds gather strength, I add to the aforesaid Composition a small Quantity of Rape Seed boil'd, without bruising, they being then able to digest that Rape Seed; but the said Rape Seed must be wash'd in cold Water, after having just boil'd up; for it nourishes the little Birds, without heating them. Now and then I give them a sweet Almond blanch'd and reduc'd to Pouder, which I mix with their Paste; and when I perceive the little Birds are somewhat heated I put in a Pinch of the Seed of Chickweed, the best grain'd and smoothest I can find. This
Com-

Composition is to be made twice a day in very hot Weather, because every thing that goes to the making of this Paste soon grows sour. This is in short what I use to bring up little *Canary-Birds* by Hand; you see it is done in a Moment. If you make use of this last Composition, you may be assur'd your Little Ones will come to good, and if you rear up 40 by Hand, scarce one will dye. If it happens that any of your little *Canary-Birds* are Sick, which is not strange in a considerable Number; instead of fair Water you may make use of the Water or Milk of Hemp, which is made after this manner. Take a Handful of Hempseed, wash it well in Spring Water, then bruise it with a Wooden Pestel, in another Water, and press out the Juice in a white Linnen Cloth, and take of it to moisten your Composition. This is call'd Hemp Water or Milk, and nourishes and warms your Sick *Canary Birds* better than fair Water, which is put to the common Compositions; but it is not to be us'd unless in Case of Necessity; because it is troublesome to make Water twice a day, and the Little Birds that are in Health do not need to be much warm'd, as the Hemp Water does. I would still add a Composition I have seen made by an Ancient curious Person, but as I know it would be thought troublesome

to

to make, and his *Canary-Birds* thrive no better than those that are rear'd with the last, I will pass it by in Silence, to avoid over burdening the Memory of those who shall read this little Treatise.

As for Provision of Biskets, it is needless to lay in any, because they are to be had at all times; only take the hardest, and those that have been longest Bak'd, which are to be given now and then to the young *Canary-Birds*,

But as for the Simnels, a Stock of them must be laid in about *Michaelmas*, when the Pastry Cooks give over making of them. You must therefore cause several Dozens of them to be made, according to the Number of *Canary-Birds* you have; string, and keep them in a dry place, and give them to your *Canary-Birds* the day you give them no Seed. They are very fond of it and it can do them no Harm. Brush the Simnel well before you give it them, because of the Dust that may stick on it.

The Simnels I cause to be made, have more Crum, and are not so flat as the others. I cause good Butter to be put into them, and a little more Salt than is usual in others, and thus they keep very well till the time when new ones are made. A little of the Crum of the Bread may also now and then be put into the Cages,
but

but it must be Stale, because the New might choak them.

CHAP. IX.

*The Hours to be carefully observ'd
for feeding of the Young Cana-
ry-Birds, when they are brought
up by Hand.*

IT is not enough to know how to make several good nourishing sorts of Paste to feed the *Canary-Birds* that are to be brought up by Hand; it is also requisite, in order to rear them without danger, to know when to give and when to refuse them their Food. If a regular course of Life be convenient, even for the strongest Body'd Men, to preserve and prolong their Life, how much more necessary must it be, without all comparifon, to rear those tender Little Birds, whom the least Excess in eating certainly kills past all retrieving; perhaps some may say, I have not observ'd any Rule of this sort, and yet I have rear'd many. But if I ask them, have any dy'd
as

as you were breeding them by Hand? they will own that many, and still more at Moulting Time. I must tell them, that the cause of their Dying whilst they are rearing by Hand, is their not being able to brook the negligent looking after them, for sometimes they are Starv'd by the long Interval between the Times of feeding them; and sometimes they are Surfeited by being fed too often and indiscreetly; and consequently Young Birds so irregularly rear'd grow Sickly, which is thought to be for want of feeding, and then the Keepers try all ways to make them open their Beaks to swallow something, but all in vain, their Stomachs being so full that they are choak'd up, for nothing digests with them, and so having pin'd a few days, at last they Dye. I have observ'd that the *Canary-Birds* rear'd by Hand, without observing any certain Rule in feeding them, are commonly so thin, poor and weak, that they are more likely to Dye than to Live, and the first Illness that seizes them, which is commonly their Moulting, they have such an ill Constitution and so infirm a Body, that they cannot withstand it, and most of them Dye. Then the Owners lay all the Fault on their Moulting, and not on the irregular way of Feeding whilst they were rear'd by Hand; and I must add, that if those
Birds

Birds happen to escape the Moulting, and are design'd to sit, the Hens often dye in Laying their first Eggs, and the Cocks being always weak and languishing, let the Hens lay addle Eggs. These reasons, which seem natural, have oblig'd me to fix a Rule for finding out the proper Hours, without being mistaken, when to feed the little *Canary-Birds* brought up by Hand, to the end that such Regularity being observ'd, they may grow as strong and healthy as if they had been rear'd by the Cock and Hen.

The Rules to be observ'd by curious Persons for feeding of their Young *Canary-Birds* are as follows, *viz.*

The first Time at half an Hour after six in the Morning, the latest.

The 2^d Time at Eight of the Clock.

The 3^d at half an Hour after Nine.

The 4th at half an Hour after Eleven.

The 5th at half an Hour after Twelve.

The 6th at Two.

The 7th at half an Hour after Three.

The 8th at Five.

The 9th at half an Hour after Six.

The 10th at Eight.

The 11th at Three Quarters after Eight, for the last Time.

Thus you see the young Birds are to be fed Eleven times a day, without any Trouble or Uneasiness, the same regular Hour being

being always observ'd. This last feeding is not always necessary, for very often the young Birds are gone to rest at that time; they must not then be disturb'd to Feed, and if it be given them it must be much less than at the other Times; for, as you may perceive there are but three Quarters of an Hour between the two last Feedings. You must make a little Stick very smooth and thin at the End, to feed them with; it must be as broad as your little Finger at most. Those who make use of a Quill cut for that purpose have more Trouble in it; because their Composition not being liquid, for the Reasons I gave in the last Chapter, the Quill bends and is not stiff enough to take up the Birds Food.

You must every Time give them their Beak full about four Times, that their Crow may not be too full, which might choak them. I must also observe, that there is more trouble in not following this rule than in keeping to it; for besides that things do not answer so well, a Man is every Moment in pain to know how long it is since they were fed, and being thus concern'd he goes to the Little Birds, who as soon as they see him do not fail to gape wide, tho' they have no occasion, for they would crave every half Hour, if they were to be regarded; then in pity you are apt to feed them, and commonly many of
them

them dye of it. I would therefore advise those who will not observe any certain Rule in this case, but their own Fancy, to leave those poor harmless Creatures to the Cock and Hen, till they are quite brought to Feed alone, rather than occasion the Death of their Birds. But as for those who are convinc'd, that there is no Pleasure without some Trouble, and who will exactly observe this Method with their *Canary-Birds*, they will perceive them to thrive visibly, and their Little Birds will by degrees grow so strong and healthy, that the Moulting, which is their most fatal Distemper, will not affect them. After 23 or 24 days, you are to forbear feeding them by Hand, especially when you observe them to pick well of themselves. As for the Lemmon Colour, and Mottled, you must continue feeding them 30 days, for they have much ado to Learn to Feed themselves. A more particular care must be taken of these Birds than of others. When they first begin to Feed alone put them into a Cage without Perches, with a little very fine small Hay, or Moss, well dry'd, in the bottom of the Cage. The first Month they feed alone you must give them what follows, *viz.* bruis'd Hemp Seed, some Yolk of Egg boil'd hard, some Simnel or Bisket very dry and grated, Water with a little fresh
Li-

Liquorice in it, a little very ripe Chick-weed; each of them apart, in the middle of the Cage, and some dry Rape Seed in their Trough. When you find them strong enough, take away all those things by degrees, and leave them nothing but their common Food, which I have spoken of before.

CHAP. X.

At what Time the Cock Canary-Birds may be known from the Hens, and the Young from the Old.

ALL the different sorts of Birds have their Marks, whereby the Cock may be known from the Hen; some are easier to be distinguish'd and others harder. For example, the Cock Linnet is easily known from the Hen, because his Wing is whiter, and he is lighter Colour'd. The Cock Gold Finch is no harder to be known, in regard the Muzzle about his Beak and his Back are very black, but the Hen has that Muzzle and the Back brown. To
spend

spend no more Time in distinguishing between the Cocks and Hens of several other sorts of Birds, I return to our *Canary-Birds*, and declare that it is harder to know the Cock from the Hen, than is generally imagin'd, especially among the high Colour'd Birds, as the Copple Crowns, the Gold Colour, and the Lemmon.

The common Rule to know the Cock *Canary-Bird* from the Hen, is his having a sort of yellow Gullet under his Beak, which hangs down considerably lower than the Hens. Besides, the Cock has a Head somewhat bigger and longer, and is commonly taller than the Hen, whose legs are shorter; and in every several sort, the Cock *Canary Bird* may be known by being of a deeper and livelier Colour than the Hen. To conclude, the last thing, which is also the surest and most to be rely'd on, for distinguishing the Cock from the Hen, is that the Cock almost as soon as he feeds alone, begins to warble; but as soon as the Moulting is over, you will hear that Cock, which before only warbled, make himself known by his singing, beyond all doubt; and his sweet wild Note, which grows louder by degrees, the next Spring at farthest attains its ultimate Perfection.

For instance, put a couple of Grey *Canary-Birds* into a Cage, and see whether what I have said, as to the knowing of
the

the two sorts, be not certain. The first glance of an Eye you cast upon the Cage, you will know the Cock, without being deceiv'd; you will perceive that sort of Yellow Gullet, a Head somewhat bigger and longer, you will commonly find the Cock taller than the Hen, and more sprightly in his Motion, and tho' Grey, he looks Yellowish by the Hen, he is so much higher colour'd. The Ash are almost as easy to be known, especially when they incline not to Gold Colour. The Colour also sufficiently distinguishes the Cripple Crowns, the Hen being perfect white, and the Cock by her inclining to Yellow. But it is very hard to distinguish between the Cocks and Hens that are high colour'd, as the Yellow inclining to Gold Colour, and the Lemmon Colour, as I have before observ'd; because the colours of those Birds being lively, the Gullet is confounded with their own Colours. Many old curious Persons are frequently mistaken in them. See what I have said about it in the 4th Chapter; nothing can then certainly distinguish the Cock from the Hen but his singing.

As for knowing Old *Canary-Birds* from the Young, it is done three several ways; first by the Colour, 2^{dly} by their Strength, and 3^{dly} by their Singing.

1. All old *Canary-Birds* have a deeper
and

and livelyer Colour in their fort than the Young; their Feet are rough and blackish, especially the Grey *Canary-Birds*; besides their Spurs are larger and longer than those of the Young ones, whose Feet are smother and their Spurs short.

2. The old *Canary Birds*, which have Moulded twice, are stronger and more vigorous, and better in Flesh than the Young, which are generally very weakly till the Spring comes on.

3. The old Birds are to be known, especially the Cocks, by their Singing, which is much Louder, and they hold it longer than the Young Cocks, who have not fully attain'd it till a Year after they are hatch'd. The same may be observ'd in the Hens, which warble louder than the Young ones, these being commonly above six Months before they open their Beaks to sing.

CHAP. XI.

At what time and in what manner Canary-Birds must be put up, to be Taught with a Flageolet.

IT is a certain Truth, that next to the Nightingale, the *Canary-Bird* sings best, and has the strongest Voice of any little Bird. When Young, they easily learn what they are taught, as Airs and Tunes on the Flageolet. For my part, I prefer them before the Nightingales, because the Nightingales are much harder to rear than the *Canary-Birds*, and besides, when a Man has so far succeeded as to breed up one, among many that dye under his Hands, He must make several sorts of Paste, as may be seen in the Chapter of Nightingales, and those are very troublesome to make, and require much Application; whereas the *Canary-Bird* is easily fed, especially when once well wean'd from the Hen. Besides, the Nightingale, after all the pains taken in feeding and rearing, generally sings but one short Season of the Year, whereas our *Canary-Birds* are continually

tinually warbling at all times. For these reasons, I think, the *Canary-Bird* ought to be preferr'd before the Nightingale, notwithstanding his Note is somewhat more Harmonious than that of the *Canary-Bird*.

As for the Time when a *Canary-Bird* is to be put into a separate Cage, in order to Teach him, that is commonly eight or ten Days after he feeds alone, and not, as many do, who take him out the first day they see he Feeds alone, without allowing him Time to gather Strength. They unmercifully shut him up in a Box, being a sort of Cage so call'd, all shut up with Boards, with some little Holes at certain Distances, that the Bird shut up there may not be stifled for want of Breath. That poor little Creature being thus scarce able to Breathe and being depriv'd of the Air, which strengthens the Body almost as much as Food, grows Melancholy and pines away, so that it often Dyes in a few Days; thus of many *Canary-Birds* shut up in that dismal Prison, scarce one escapes. If you desire to succeed better in that Point, you must observe this Method.

A Fortnight, at soonest, after your *Canary-Bird* has been taken from the Hen, if you perceive he begins to warble, which is a Sign that he is a Cock, and at the same time in good Health; then part him

D

from

50 *A New Treatise of*

from the rest, and put him into a Cage cover'd with a very thin transparent linnen Cloth, for the first eight Days. Put him into a Room distant from all other Birds whatsoever, so that he may never hear any of their Wild Notes, and then play to him upon a little Flageolet, the Notes whereof are not too high, for if your Flageolet is too loud, your *Canary-Bird*, who will not fail to follow the same Tone, when he has learnt the Air, and will repeat it several times in a Day, will so dry up his Lungs, that it will cause him to grow lean and Dye in a few days.

At the end of a Fortnight, take away that thin Linnen Cloth that covers his Cage, and supply it with a Piece of thick green or red Serge, and leave him still in the same Place, till he perfectly learns what you teach him. Some *Canary-Birds* are much more apt to learn than others; for some are perfect in two Months, and others not in six, but you must have Patience, when you resolve to teach them any thing, for without it nothing can be done. When you give him his Food, which must be enough for two Days at least, it must be done by Candle and not by Day Light, that he may not be distracted, but may learn what he is taught the sooner.

As for the Tunes, he must be taught only one fine Prelude and a choice Air; when

Canary-Birds.

51

when they are taught more, they are apt to confound them, and being taught too much, they often know nothing perfectly, their Memory is so over burden'd, that they know not what they sing. Besides their Distemper of Moulting, which is once a Year, and lasts above two Months, diverts them so that they do not sing, and consequently easily forget the several Tunes they have been taught with so much trouble.

Do not follow the Notion of those, who think that the many Lessons they give their *Canary-Birds* every Day, improves them the more; they tire themselves and only make the Birds uneasy; five or six Lessons a Day are sufficient for a *Canary-Bird*, tho' he be never so little dispos'd to learn; besides the trouble would much outweigh the Pleasure, if a Man were to be continually about his *Canary-Birds*. You must therefore give them but two Lessons in the Morning when you rise, another about Noon, and the same before you go to Bed. The Morning and Evening Lessons do the Bird more Benefit than all the rest during the whole Day, because at that time when all is still, the Birds being less diverted, better retain what is taught them, the Tunes taught them must be repeated 9 or 10 times at every Lesson, and those same Airs must be play'd running,

running, without repeating the first part of them twice, and the Conclusion in the same manner, as is usually done in Concerts, or other Playing. A little Prelude may be given them in *C Sol Ut*, which all Men know is made for Birds, with a March, as that of the Guards, or any other, which is also in *C Sol Ut*, as well as the Prelude. When a *Canary-Bird* has learnt those two little Pieces perfectly, every Man ought to rest satisfy'd, without taking the trouble to go about Teaching them any more. They that would meet with good Success, must not undertake to teach *Copple Crown*, nor *Lemmon Canary-Birds*, which have not strength to overcome the Diseases incident to them, nor so strong a Voice as the Grey. A *Grey Canary-Bird* of a good Breed, with a white Tail, is more apt to learn by a *Flageolet*, than any other sort of *Canary-Bird*. You must not pretend to teach two *Canary-Birds* in the same Room, much less in the same Cage, as many have attempted to do. If that be done, it must be only for a Time, that is, as soon as one of the two begins to take the Tune, they must be immediately parted at such a distance, that they may not hear one another, because they will do nothing but distract one another; and it would be all Time lost to them and to him that teaches them

them. This may suffice as to this Subject, let us proceed to another.

I have thought fit to prick down the Prelude and Air I have spoken of.

CHAP. XII.

Of the several Dispositions and Inclinations of Canary-Birds.

IT may truly be said, that almost all *Canary-Birds* differ in their Temper and Inclinations; but being afraid it would take up too much Time to distinguish between them all successively, I think fit to divide them all into four Classes.

In the first Place, I find some Cocks who are always of a Melancholy Temper, and, if I may so call it, thoughty, who sing but seldom and in a doleful Tone, being for the most part huffed up. This sort of *Canary-Birds* seldom answer the end they are design'd for; for if you would teach them to sing Flageolet Tunes, they are a prodigious Time learning, and are never perfect in what they have been taught; besides, that what they have learnt they easily forget, especially the first time

they come to be Sick, as at Moulting Time, or the like; and sometimes there are such as pine so much at being always cover'd, that they Dye; and this sort of *Canary-Birds*, tho' they are Cocks, I believe would never Sing, unless they were put to it under other old brisk *Canary-Birds*, that hearing them continually Sing, they may in some manner serve them instead of Masters. These same Birds are sometimes naturally so slovenly, that their Feet and Tails are always very Dirty; when you would clean them without Danger, you must use the following Precautions.

Take the *Canary-Birds* into your Hand, and with a little Spittle by degrees, take off the hardned Filth that sticks on their Feet, which often hinders them from perching; and it often happens, that when care is not taken to clean them, their Feet grow sore and their Claws drop off. If you will make use of Water instead of Spittle, to clean them, it must not be cold, unless it be in very hot Weather, for besides that the cold Water does not clean them so well, there is Danger that it may kill them, that Coldness striking to them on a sudden, especially in Winter. Some young curious Persons have lost their *Canary-Birds* by using of cold Water, at a Time that did not seem to them Cold enough to warm it; and your Hands ought
to

to be hot when you take hold of the Bird.

All the particulars I take Notice of in this small Treatise, are of such Consequence, that those who through Ignorance or Negligence do not observe them, loose many of their *Canary-Birds*; for those little Creatures are of so tender a Constitution, that a very small matter impairs their Health and occasions their Death.

On the other hand, if you design to have this sort of *Canary-Birds* to Sit, there is not only Reason to believe that their Young Ones will be no better than the Old, but the first cross Accident that happens in their breeding, as the Death of any one of the Little Ones, or any Distemper of the Hen, or any other thing of this sort, they grow so melancholy and dull, that they often Dye, or if they happen to survive it, they are all the rest of the Time they remain in their Huts, so heavy, that they never hear the Hens by Singing, especially when they perceive the young begin to Hatch. I would advise those who have any of these, to get rid of them, to save themselves the Concern it will be to them, to see that whatsoever they can do for them, nothing will succeed.

There are other Cock *Canary-Birds* so mischievous, that they kill the Hens that are given them to Tread; but sometimes

it happens that those Cocks have some other Qualities which make some Amends for that Fault; as for Instance, they Sing very sweetly, have very fine Feathers, and are very familiar; therefore there is no need of getting rid of them, since they have good Qualities to recompence the bad. Besides, if, for want of another Cock you are resolv'd to have a Breed from them, you may do it as I will here prescribe.

Take two very strong Hens, and, if possible, let them be a Year older than that mischievous Cock you design for them; put these two Hens sometimes together into the same Cage, that they may grow very well acquainted, and by that means not grow Jealous and Fight, when they have but one Cock between them. A month before they are put up in order to sit, they are both to be turn'd into the same Hut, and when the proper time for coupling is come, put in the Cock to the two Hens; he will be sure to beat them, especially during the first days he is with them; but both the Hens joining to defend themselves against him, they will at length gain an absolute Ascendant over him; so that perceiving he can gain nothing upon them by Force, he will in a little time grow so familiar with the two Hens, as to gain them by Kindness; and thus these
Weddings

Weddings by Force often prove more successful than others, from which great Hopes were conceiv'd, and in the end come to nothing, or very little. I have taught this way of coupling mischievous *Canary-Birds* to some, who have own'd that it has prov'd successful.

There are others of so base a Temper; that they break and eat the Eggs, when the Hens have laid, and if they happen to let the Hens hatch any, no sooner do the Little Ones peep out, than they cruelly seize them with their Beaks and drag them about the Hut till they dye. The Method to prevent this Accident, and to make that sort of Cocks fit, is to shut up the Cock, the Day before the Little Ones are to be hatch'd, in a little Cage, in the middle of the Hut, and there give him Meat and Drink. There seeing the Hen go backward and forward, he will not fret, and the Hen, if she is good, will rear the Young Ones very well alone, without his Help; at the end of 10 or 12 days you must take away her Little Ones to bring them up by Hand, that she may not spend herself too much. As soon as the Little Ones are taken away, be sure to turn the Cock loose to her, and every time she Sits, do as has been said above. But if he happens to break and eat the Eggs, that is harder to be remedy'd, because he cannot

58. *A New Treatise of*

be shut up whilst the Hen is laying, because her Eggs would prove addle. However this Remedy may be apply'd; as soon as the Hen has lay'd her Egg, that very Moment, lest the Cock should eat it, you must take it out of the Nest and put it into a Deal Box, full of the Sand us'd for making of Glasse; that Sand is very fine; the Egg is in no Danger of breaking on it, and always keeps fresh; if you will, you may cover the Egg with Moss and shut up the Box, which put into a Place where no Body may meddle with it. This I have here observ'd is to be understood for putting in the Eggs of all the *Canary-Birds* you have. The Box is to be made purposely for that use, and to have as many Partitions as you have Couples of *Canary-Birds*. To avoid mixing the Eggs that belong to one Couple of *Canary-Birds* with those that belong to another Pair, the Name of the Pair they come from is writ on the cover of the Box, so that it may be exactly over the Partition the Eggs belong to when the cover is on, and thus you easily know to which Pair every parcel of Eggs belongs. But to return to the mischievous Cock, you are to take from him the first Egg his Hen lays, and lay an Ivory one in the place, do the same the next day; always taking it away the Moment the Hen lays, that the

the Cock may not have time to break it. When the Hen has laid her last Egg, she being no longer in need of the Cock, shut him up in a Cage as was said before, and give the Hen the Eggs to Sit on. The Cock is to remain in his Cage, in the middle of the Hut all the time the Hen continues Sitting, and rearing her Young; but as soon as ever you have taken away the Little Ones to bring them up by Hand, there being no more Danger, you must let out the Cock again to the Hen. I know, what I have here prescrib'd in Order to prevent ill Accidents, is somewhat of a Confinement; but I only propose it to such as have Patience enough, and are resolv'd to breed from such ill natur'd *Canary-Birds*.

Lastly, those I assign to the 4th Class, are *Canary-Birds* that are always Merry, their Throats being continually open, singing long and often. They are so Tame, that they take whatsoever is given them, tho' it be out of a Mans Mouth. These may be call'd perfect *Canary-Birds*, being fit for any thing they are design'd; for if you will have them breed, they please the Hen with their harmonious Singing, which revives her very much, especially whilst she is sitting. These Birds are so good natur'd, that they will often Sit upon their Hens Eggs some Hours in a day, and feed the

the Little Ones as soon as hatch'd, to ease the Hens. On the other Hand, if you have no mind to have these Birds breed, but intend to teach them some Flageolet Tunes, they will learn sooner, and sing louder than others. When you happen to light upon such good Temper'd *Canary-Birds*, never think of changing them for others of more beautiful Colours. These Birds, tho' they be but plain Grey, are to be preferr'd before Copple Crowns, who have no other good quality but their beautiful Feathers. There are some curious Persons who when they meet with any of these good *Canary-Birds*, are willing to take them in exchange, tho' Grey, for Copple Crowns, as well knowing that those Grey Ones, will be worth more to them, by reason of the Number of Young Ones they will Hatch and Rear, without fearing they should Dye.

CHAP.

CHAP. XIII.

*Of ill Accidents Canary-Birds
are lyable to in sitting.*

BEfides the cross Accidents that may be avoided, which happen to *Canary-Birds*, when they are in Huts, whereof I have spoken in the last Chapter, there are others befalling, notwithstanding the good Disposition of the *Canary-Birds* that are put into Huts, and all the Care taken of them.

For Instance, a *Canary-Bird*, tho' he have all imaginable Accomplishments, is not exempt from falling Sick, and that often, when the Hen stands most in need of him, as when she is about laying, or when the little ones are 7 or 8 Days old, which is the Time when a good Cock *Canary-Bird* is to ease his Mate of the Burden of feeding them, that she may rest, in order to go about making another Nest immediately, to Lay in it again. If then the Cock is taken ill at that Time, a Man is hard put to it, and knows not which way to turn himself, especially if he has but newly undertaken to breed *Canary-Birds*.

But

But since, as is commonly said, there is a Remedy for all Things but Death, the Method to be us'd in such an unlucky Case is this.

Take your Sick *Canary-Bird* immediately, and put him into a little Cage; then endeavour as near as you can, to find out what Distemper it is that has seiz'd him, and having discover'd it, apply the proper Remedies with all speed; I shall speak of them in another place, when I come to discourse of the several Diseases *Canary-Birds* are subject to. Put the Sick Bird into the Sun, spurt a little white Wine upon his Body, which is a Remedy that agrees with all their Distempers, after which apply what other Remedies you shall think fit. If they have no good Effect, but on the contrary you perceive your Bird grows worse, and that the Hen begins to pine much for the Absence of her Cock, it is then Time to think of providing another Cock, to supply the Place of the Sick one; but you must not follow the Opinion of those, who think that a Hen cannot be several Days without the Cock. I have seen some that have rear'd their young ones very well, tho' the Cock were Dead. It is true there are others not so indifferent, but 8 or 10 Days may pass very well without any Danger of the Hens dying for want of the Cocks Company.

pany. You may also show her the Sick Cock, putting of him into the middle of her Hut, in a little Cage, for some Hours; but this is only to be done when you see the Hen very uneasy and restless, for want of her Cock.

Without seeing the Sick Cock I can tell, that his Sicknes proceeds, either from having heated himself too much with his Hen, or from having eaten too much of nourishing Food, such as must be given them when they have little ones to rear.

The infallible Remedy for the first of those Distempers is 8 or 10 Days Rest; and for the Second to Diet them for some days, in order to bring them down, giving them no other Food but Rapeseed. A few Days after you have made use of these Remedies, put your Bird again to his Hen, who will be as merry and well pleas'd as before; but if he falls sick again, he must be taken out and return thither no more, tho' he recovers; for it is a certain Token that he is too tender to endure the Trouble and Uneasiness of bringing up the Young.

What I have said of the Cock Canary-Bird, is also to be understood of the Hen, only with this Exception, that when the Hen falls sick, whilst she is Sitting on Eggs, those Eggs must be taken out of the Hut,

smol

at

at the same time with her, and lay'd under other Hens that have been sitting about the same Time. If she falls Sick : when the little ones are hatch'd, you must consider whether they are strong enough to rear by Hand, and if not, they must be given to a Hen that has young ones of about the same Age, for her to feed them. It may happen that the Cock will feed the little ones, without the help of the Hen; however they must not be left to him, especially if they be but 5 or 6 days old; for at that time they have as much need of being sat upon, as of Feeding, because they are not fledg'd; and tho' the Cock keeps them from dying for Hunger, they will be starv'd with Cold, for want of the Hen.

There are Accidents that happen for want of Precaution, as the breaking of Eggs through Heedlessness. For Example, a Hen lays her Egg very early in the Morning, in some close Corner of her Hut, you come in the Morning to remove the Sand, in Order to make it clean, and find too late, that you have broke an Egg, and perhaps it is the loss of a *Canary-Bird* that might have been worth 2 Guineas, which would have come from that Egg. When therefore you do not find the Egg you expected the Night before, in the Nest, you must rather look about every Nook of the Hut for it, then feel with your Hands; I have often found

some among Chickweed and Groundsel we are oblig'd to give the old ones, especially when they have Young. Having found the Egg, take it up gingerly by the two ends, for that way it is less apt to break than if taken up by the middle, and lay it in the Box I have before spoken of.

There is a Distemper the Hens are sometimes subject to in a few Days after they are in their Huts. In a moment you see the Hen all huff'd up, and refusing to eat, and sometimes she is so very ill, that she cannot stand upon her Feet, she lyes down on the Sand, and unless speedily assisted, will soon breathe her last. This Distemper commonly seizes them towards the Evening or in the Morning early. In this case you must take the Sick Hen in your Hand, and being well assur'd that her Distemper is occasion'd by Laying, you must, with the Head of a great Pin, put some Oil of sweet Almonds up her Fundament, which will dilate her Pores, and she will afterwards lay her Egg with ease. If you perceive she continues ill, you may make her swallow a Drop of that same Oil, which will ease the Gripes and sharp Pains she endures. You must leave her in a little Cage, with some fine Hay, place her in the Sun, or before the Fire, if the Sun does not shine, till she is restor'd to her former

former Vigour. Give her good Food, as boil'd Seed, Bisket, and dry Simnel, Pink-feed, and if you perceive, that notwithstanding all that good Nourishment, she can scarce recover, spurt some little White Wine upon her with your Mouth, and make her swallow a little luke warm, with a little Sugar-Candy, or other. These things will restore her to her former Health. This Distemper seldom seizes them but at laying their First or Second Egg.

It also sometimes happens, that a Hen pulls the Feathers off the Young Ones, when they first begin to shoot out, which is commonly when the little ones are 7 or 8 Days old. There are two several ways to obviate this Accident; the first to take away the Young Ones, if they are strong enough to be rear'd by Hand; the Second is, to put the Little Ones, if you must leave them, into a little Cage, with their Nest plac'd in the middle of it; the sticks of the little Cage must be at a good distance from each other, and then the Cock and the Hen wil feed the Little Ones between those sticks, without plucking their Feathers.

It also happens that the Hen sweats over her Young, when they are but 2 or 3 days old, and sometimes as soon as they are hatch'd. This Distemper is discover'd by the
the

the Feathers under the Hens Stomach and Belly being wet, and the hard coming of the Down on the Little Ones. When the Young are six days old, and the Hen begins to sweat, they are then out of Danger; but many Dye before they come to that Age, being stifled by that Distemper. I will hereafter speak of several Remedies that may be us'd at that time; but in regard that all the Remedies I shall prescribe against this Distemper, are very difficult and hard to prepare, besides that the care is not always certain, the greatest and most infallible Remedy, is immediately to take away the Little Ones from under the Hen, and, if you have not any yourself, to find out some Friend who has Young *Canary-Birds* of about the same Age, to put them under his Hen, with hers, to be rear'd. These small Courtesies are reciprocal among curious Persons.

It often happens that Hens lay 3 or 4 Eggs at their first breeding, and then leave them; when that happens, after they have lain 2 or 3 days in the Nest, to try whether she will not bethink herself to sit on them, and you perceive that she does not go to them in that time; but that on the contrary she spoils the Nest the Eggs are in, you must take them away, and lay them under other sitting Hens. I have observ'd that the Eggs those Hens would not sit up-
on

on were generally addle; for some *Canary-Birds* are sensible when their own Eggs are addle, and then they will not sit upon them. I have also put other addle Eggs under some Hens, instead of their own, which they broke and threw out of their Nest, almost as soon as I had laid them in, and I was fain to give them counterfeit Ivory Eggs to please them till they had done laying, when they are to have their own Eggs given them, that they may begin to sit upon them all the same day. You must not be discourag'd when you see a Hen leave her Eggs at the first sitting time; that most frequently happens to - Young Hens that have never sate before; for the same Hens perform Wonders at all the following times of Laying, continually Sitting on their Eggs, and rearing their Young Ones very well. However, there are some Hens that will never Sit, or only the last time of laying; such Hens are rarely met with; besides those who have any such, may let the Hens lay, and give their Eggs to others to sit on, after having left the Eggs of every time, a Day or two in the Nest, to try whether they will not come to them.

It is not to be wonder'd at, if there happen to be some Eggs short of what there was in the Nest, even after they have been several days sate on, the reason of it is that

that the Hen having perceiv'd them to be addle, has taken them out of her Nest and eaten them; the very Shells of them are not often to be found. Many who do not understand this, would be apt to suspect somebody might have taken them away, but without any reason.

Another Accident sometimes happens unforeseen, which is to find a *Canary-Birds* Foot broken in the Hut, and some do not know how that comes about, and the reason commonly is, because the holes made in the Elder Sticks are too big, and the Birds Claws too long, so that the Birds sometimes thrust their Claws so far into those holes, that they cannot get them out again, and if you do not happen to be there that very Moment, they flutter so violently, that they seldom get off without breaking their Toes. To prevent this accident two things are to be done; First, the holes in the Elder Sticks must be made no bigger than the point of a Needle; Secondly, the *Canary-Birds* must never be put into Huts, without looking first whether their Claws are too long, which is frequent in old *Canary Birds*, and then you must not stick to cut off about half the Length of them at most, for should you cut them too short, they would not be able to stand upon their Perches.

Special care must be taken that the Perches in the Hut be well fix'd that they may not fall, especially when the Cock is going to tread the Hen, for it would occasion her to lay many addle Eggs.

There are other troublesom Accidents, which I will not omit, being such as *Canary-Birds* whose Claws are too long are subject to. The Hen very often does so entangle her Claws in the Nest, when she would go out, that she commonly drags the Nest after her and breaks the Eggs, or kills the Little Ones, if they are hatch'd.

Sometimes the Hens Claws are so sharp, that they strike through one of the Eggs she sits on, which taking Air, can never come to good.

Sometimes as she comes to sit over her Little Ones new hatch'd, her Feet slip upon the Body of some One of them, which being prick'd with her Claws, Dyes of it.

Another unlucky Accident that may happen, is when you think a Hen feeds her Little Ones well and she does nothing but sit on them; when you have carefully observ'd for half a day at most that she carries nothing to the Young Ones, you must without losing more time, take them away from her, and lay them immediately under another Hen, that feeds them well and that has

has Little Ones of her own near about the same strength as those then given her. If you happen in one Nest to have one Little One weaker than the rest, and some others like it in another Nest, change them, laying the strongest together and the weakest by themselves, not leaving above 5 or 6 to a Hen at most; and she must be a very good Breeder to rear so many, without losing any.

When you have any Hens which you suspect do not feed their Little Ones well, as is frequent among the Mottled and White Ones with Red Eyes, as also the Ash and Lemmon, and lastly some Copple Crowns, whom you know by Experience not to have been good Feeders, you must before they are hatch'd put them under Greys, which the curious call Nurses, throwing away the Eggs of those Greys, if you have not other Hens to lay them under. It is sufficient that a Hen has fate 4 or 5 days to give her Eggs that are ready to hatch. Those who live in the Country may lay their *Canary-Birds* Eggs into Gold-finches Nests, and they will be sure to have Young ones without any Trouble, provided they take care not to put in Eggs that are unfit, that is, not to put *Canary-Birds* Eggs which have not been fate on, into a Gold-finches Nest, whose Eggs are very forward; or the contrary,

trary, in both which cases they will come to nothing. Therefore when you have found a Gold Finches Nest you are first to break one Egg, and by that you perceive whether they are advanc'd, that you may put in *Canary Birds* Eggs that have been late on about the same time.

When the Little Ones are 10 or 12 days old, they are to be taken away and rear'd by Hand; if you will have them continue longer you must put them into a low Cage, with a little Net over it, that when the Cock and Hen come to feed them, they may receive their Sustenance. When they have been fed some days, you may by degrees bring them nearer to the House, always setting the Cage in a very open Place; and when the Little Ones come out of their Nest, you must put them into a larger Cage and leave them in the same Place, till you perceive that the Cock and Hen go no more to it. During all that time you must put something to eat into their Cage, as the Yolk of an Egg boil'd hard, and bruis'd Hempseed, that they may learn to eat alone, tho' the old Ones feed them at the same time.

No Nests of any other Birds are proper for them. The same might be done in Linnets Nests; but that they commonly forsake their Eggs, when they perceive they have been meddled with.

The

The Yellow Hammers disgorge; but they feed on a sort of Seed which kills *Canary-Birds*, and therefore you must not venture to give them any to rear in their Nests, unless it be when they are coupled with *Canary-Birds* in Huts to breed Mongrels, where at that time they have no Food given them but what is proper for *Canary-Birds*.

When a Hen happens to fall Sick, some days after her Eggs are Hatch'd, or that she forsakes them, as sometimes has been seen, you must then, for want of other Hens to put them to, buy a Nest of young Sparrows, and put as many of them as is convenient into the Nest of your forsaken Birds, that being altogether they may keep the little ones warm, and then you must Feed the little *Canary-Birds* every Hour, till they are twelve days old, after which you must feed them as prescrib'd in the Ninth Chapter. If the Weather be somewhat cold, lay over the Nest a piece of very fine Lambs Skin. You may feed the Sparrows with more common Food than that of the *Canary Birds*, that they may not grow big too fast. A Nest of Linnets might be made use of, but they are not so hot as Young Sparrows. By these means I have sav'd some, which would certainly have perish'd in other Hands, where this Precaution had not been us'd.

E

These

These are the most common Accidents that may befall *Canary-Birds* when they are in their Huts.

C H A P. XIV.

The manner of putting several Hen Canary-Birds to one Cock, in Huts, or Voleriers.

IF you have more Hens than Cocks, and will not be at the Expence of buying others, who may make them breed as follows.

When a *Canary-Bird* is strong and vigorous, which is known by several Tokens, as when he Sings very loud, long, and often in a day, and is so brisk, that he cannot stand still one moment in any part of his Cage or Hut, you may, without any difficulty, give him Two Hens.

To this purpose you must place two little Huts side by side, each of them having a Door of Communication near one another: Put the Two Hens apart into the Two Huts, and turn the Cock into one of them. The Cock being call'd by both Hens will go sometimes to one and sometimes

to the other, and by that means Tread them both, which will spare you one Cock.

Here follows another Method.

If you have but one Cock, as above qualify'd, and but one Hut, provided it be pretty large, you may also put your Two Hens into it with the Cock, making a little Partition in it with a Board, that the Two Hens who are in their Baskets at the ends of the Hut, may not be distracted by looking upon each other. The little Board, I have mention'd, must be very thin, and at most not reach above a Quarter of the height of the Hut, that being enough to hinder the Hens seeing one another, when they are Sitting on their Eggs. That Partition reaching no lower, it will be easier for the Cock to carry Food to the Hens. That all things may answer the better, the Hens must have been kept together some Months before, to the end they be well acquainted and us'd to one another, when you put in the Cock you have design'd for them; and if it should happen, that when they have both lay'd, they should not Sit, as being distracted by being Two Hens in one Hut, you must then take away the Eggs and give them to others to Sit on. The Eggs those Two Hens lay will be as good as if each of them had a Cock apart. Thus you will save a Cock Canary-Bird and a

Hut. Experience will show that this is very easily done, and beneficial for those who practise it. Sometimes there is a necessity for so doing, when the Cock happens to be so sprightly, that he grows uneasy and restless, when the Hen leaves him to Sit upon her Eggs. Then having another Hen with him, he is pleas'd, going sometimes to the one and sometimes to the other, which often prevents his breaking the Eggs, or killing the Young Ones, which he does to divert himself, or in Spight, because he finds the Hen indifferent towards him, and too regardful of her Eggs or Little Ones.

There is still another way of putting several Hens, with a small Number of Cocks, into a little Closet or Volery, very light, and if possible, expos'd to the Rising Sun. Take all things out of it, during the Four Sitting Months, and fill it with *Canary-Bird* Cocks and Hens. You may turn in at least twice as many Hens as Cocks, that is, if you turn in 12 Cocks you may give them 24 or 25 Hens; you may place in several parts, as many little Baskets as there are Hens, and in the middle of the Closet lay all that is requisite for making of their Nests. In the middle of the Closet must also be a Table and on it three or four great Troughs full of Water and their common Seeds. If there should

should be but one Trough, so many Birds could not Feed all at once, and they would often Fight to come at the Meat first. You must also put in long Elder Sticks for them to Perch on. There must be a Wire Grate, that the Glafs Windows may be open'd, when the Weather is Fair, to give them the Air, and that they may not fly away. Every Hen will make her own Nest, and never mistake or go into another's. You may place about the Closet some Boxes with Greens, as Orange Trees, or the like, which makes them chearful, and several Hens may make their Nests there, if a Basket be plac'd in the middle of a Box. Those who have the Conveniency of a little Closet, have much more Pleasure and less Trouble than others. They see their Birds fly about, as if they were in the open Country. The Eggs may be also left in every Basket, without taking them away, as has been said before; and thus you will be at no other Trouble but giving them Meat for several days, and keeping the Place extraordinary clean to prevent the Mice getting in, who would kill the Cocks and Hens and devour the Little Ones.

The Young Ones are to be taken out of their Nests after 10 or 12 days, if you design to bring them up by Hand; if not, leave them with the Cocks, giving

the Hen another Basket to make a new Nest.

CHAP. XV.

How many Eggs a Hen may lay in a Year ; and how they may be known to be good.

IT is to be observ'd, that almost all Hen *Canary Birds* differ from each other in laying.

Some do not lay at all, and those are call'd Barren Hens.

Others are so ill Breeders, that they lay Eggs but once, or twice in a Year ; besides, after they have laid their first Egg, they often rest the next day, and do not lay the next till two or three days after.

Others lay only three times, and are regular in each time of laying, having 3 Eggs at every sitting, without any days Intermission.

There is a 4th sort, which may be call'd Common, being very numerous ; they Sir four times, and every time lay four or five Eggs ; but they are not always regular in laying.

There

There are still others that have more Eggs than all those I have mention'd; they Sit five times, and would do it oftener if they were permitted; every time they lay 6 or 7 Eggs. When this last sort of *Canary-Birds* feed well, they are excellent, too much cannot be made of them; they are worth more than six others of the common sort.

If you would know how many Eggs Hen *Canary-Birds* may lay in a Year, I desire you to look back upon what has been said, and you will at once perceive that

There is no reckoning the Eggs of the first sort, because they lay none.

Those of the 2^d sort may amount to 4 or 5.

Of the 3^d sort there may be 9 or 10.

Of the 4th sort there may be about 18.

And lastly, the Eggs of the last sort, which, as I have observ'd is extraordinary, cannot well be ascertain'd. I shall only take Notice, that if you let them go on Laying, that is, till the Moulting time, they will in a Year lay 35 Eggs at least. As for the knowing of the Eggs it is easily learnt.

Some say they know whether an Egg is good the very moment the Hen has Lay'd it, before she has Sate upon it, which is false. They weigh it in their Hand, and

find it, as they say, somewhat heavier than another Egg they suppose to be Addle; and upon this Conceit they decide of the goodness of the Egg. If a Man were to lay a Wager upon that Trial only, he would be as likely to Lose as to Win, because it is meer Chance, and not Knowledge that determines; you may as well affirm that the Eggs are good as soon as they come from the Hen, without having been then sate upon.

When you have some Pairs of *Canary-Birds*, from whom you have found no Addle Eggs to come, and other Pairs on the contrary, which have never produc'd any good, you may be almost sure to win any Wager you lay upon them, and those who shall happen to lay Wagers with such curious Persons, will be fully perswaded, that it is the Weight they find in the Eggs which makes them distinguish between the good and the bad, and yet you see they would be grossly mistaken.

Do not therefore depend upon that gross way of distinguishing, which may often occasion you to throw away an Egg you suppose to be Addle, which would have been good, if sate upon.

To avoid being deceiv'd, you must follow the common Rule, which is to look upon the Eggs, when the Hen has sate upon

on

on them '6 or 7 days. Take the Eggs then from under the Hen, and view them against a Candle, or the Sun. If you perceive that those Eggs are thick and grow heavy, it is a sign that the Little Ones are forming in them, and that they are good; if on the contrary you find they continue as clear as they were the first day you gave them to the Hen to Sit on, it is a certain sign that they are Addle, and then, especially if you examine them the 7th day after the Hen has fate on them, you may without any scruple throw them away, for they will only serve to tire the Hen to no purpose.

If you have several Pairs of *Canary-Birds*, whose Hens have lay'd near about the same days, you may take away the Addle Eggs from every Hen, and of three leave only two Sitting, giving 5 or 6 Eggs to an able strong Hen, because very often some one of the Little Ones dyes in the Shell, and receives no nourishment; and by that means you bring the Third Hen, whose Eggs have been given to the other two, to fall to work about making another Nest.

When your Hens have laid their first Egg, it must be taken away immediately, and an Ivory one put in the place of it, to amuse them. You must not put in old Addle Eggs, as most Men do, which often break in the Nest, and infect it; and the

ill Scent may make the Old Ones Sick; take away also the second Egg, and add a second of Ivory, and do the same at the third and fourth. When you perceive the Hen is like to lay no more, return her very early in the Morning her own Eggs, taking away the false ones of Ivory. Do this at every sitting, for should you leave the Hens their Eggs, and not take them away, they would Hatch at several times, and the first Young Birds Hatch'd, being stronger than those which would come two days after, would take all the Food from the Hen, and would stifle the last comers, or scratch and disturb them with their Claws.

A Hen must have laid her Egg by 7 of the Clock in the Morning, or 8 at the latest. If she stays longer, it is because she is Sick, and if you perceive it, she must then be help'd, as has been said before.

Some are apt to admire, that of 4 or 5 good Eggs, found to be good, by looking through them on the 7th day the Hen had been Sitting on them, very often only two Hatch. That Accident may proceed from several Causes, the first is sometimes the often handling of the Eggs whilst the Hen is Sitting, which cools them so much, that the Young Ones in them take no Nourishment, and that hinders their thriving; or else it is because those Eggs being so often

often handled get some little Crack, which is inperceptible, and if never so little Air gets into the Egg, the Little One in it Dyes immediately. This frequently happens to those who are but newly us'd to *Canary Birds*, for they being Novices, think every thing so fine and wonderful, that their Hands are as busy as their Eyes. To avoid this Misfortune, the Eggs must be touch'd but once, that is, the Day when the Addle ones are to be taken away, leaving the Hen and her Eggs to themselves after that.

This Accident may also be occasion'd by the Hen, who having too many Eggs to Sit on, by changing their Places very often, sometimes thrusts one from under her, which grows so cold, and remains there so long unobserv'd, being hid under the end of her Wing, that the Young One in it takes no more Sustenance, and consequently comes not to good.

This is the only Occasion you must take to touch the Eggs, putting that you find cold in the middle of the others.

Thunder is another Accident; for when there happens any extraordinary Thunder on the 7th or 8th day after the Hen has been Sitting, the Little Ones which are scarce half form'd, sometimes perish, and the Eggs that were thought to be good, because they were dark and somewhat hea-

vy,

vy, are quite spoil'd by the Thunder. When the Thunder happens in the day time and the Hen is gone off the Eggs to ease herself or feed, there is much cause to fear the Eggs will be spoil'd, and therefore you must then endeavour to make the Hen return to her Nest; for, when on them, she commonly saves them from that Accident; and therefore the Thunder is not to be so much dreaded at Night, because the Hen is then on her Eggs.

Some put a bit of Iron into the Nest, to hinder the effect of the Thunder; but I will not prescribe that as a sovereign Remedy. When the Thunder happens on the 11th or 12th day the Hen has been Sitting, it is not so much to be fear'd; because the Little Ones are stronger, tho' still in the Shell, and very often it only makes them Hatch 24 Hours before their usual time.

However, Thunder sometimes is so violent that it destroys both the Old and Young, and if they are never so little out of order otherwise, that Thunder makes an end of them, and they are found dead and stiff in their Huts, without any other Cause.

The Eggs are generally 13 times 24 Hours under the Hens; for Instance, you lay 5 or 6 Eggs under your Hen on a Saturday at 7 in the Morning, and you will have

have Young Ones hatch'd the 2d Friday following in the Morning. Some, but very rarely, Hatch one day before the usual time; either the Thunder, as was said above, or the great Heats, as in *July* and *August*, may cause them to advance; others on the contrary, come 24 Hours later, or thereabouts, and that is occasion'd by Cold, as sometimes happens at the first Sitting in *April*; or it may proceed from having handled the Eggs too often, as has been said; or lastly, the Hens being Sickly or Weaker than others, the Eggs under them may Hatch a day or more later than usual.

The Fear of breaking the Eggs when handled, makes us take them up trembling, and even so we often break them, either pressing them too hard between the Fingers, or else letting them fall for want of holding fast. To obviate this Accident, which daily happens to the most experienc'd Persons, they are not to be handled unless in case of Necessity, and when you cannot avoid it. You must take them up unconcern'd and without Dread, for the Fear of breaking them often proves a dangerous Precaution. If therefore you boldly take up your Eggs in your Fingers by the ends, but never by the middle, as I observ'd before, you will be sure not to break them; and if still you be apprehensive, notwithstanding

standing all these Precautions, you may make use of a little Coffee Spoon that is very thin, to take them up with, and that way you will still be safer.

CHAP. XVI.

At what time a Hen Canary-Bird toils most, whether when she Lays, when she Sits, or when she Rears.

EVERY experienc'd Person speaks variously as to these several Articles; some say the Hen is more fatigu'd with Laying than at any other time; for, say they, Nature must at that time make some extraordinary Effort, which is the reason that many Hens wanting Strength to bear the Pains they endure at that time, often Dye, notwithstanding all the Remedies apply'd to them. Others assert, that a Hen, which is 24 or 25 days sitting on her Eggs and on the Little Ones after they are Hatch'd, suffers more than in Laying. A Hen, say they, that is so long without going out of her Nest, unless it be now and then a Moment

Moment in haste, during the day, upon such pressing Necessities as cannot be avoided, as particularly to eat and drink, when the Cock does not bring her enough, is heated, and endures more than a Hen that has but an Hours Pain and sometimes less, when she is to Lay her Egg; whereas this other, lying continually Day and Night in the same Posture for near a Month, very often grows so Sickly, that she is sometimes found in her Nest with her Head under her Wing, and ready to dye, unless she be immediately taken away to rest some time, giving her Eggs or Little Ones to another to look too, that is either stronger, or less fatigu'd,

For my Part, I affirm, that the Hen which is rearing her Little Ones suffers much more than that which Lays or Sits; because the Laying Hen suffers not above an Hour at most, as has been just said; the Sitting Hen is often inur'd to that quiet Posture she is in, and the Cock commonly bringing her Food to the Nest, eases her of all care, so that very often she grows so Fat, being satisfy'd with her Condition, that it is troublesom to bring her down again. In short, she is so well pleas'd, that when a Man goes about to make her leave her Nest, she is for pecking at all that comes near her, to shew she is well satisfy'd there, and unwilling to

to remove; but this Hen tires herself with feeding her Young, and very often the Cock, who carries her no Meat, leaves all the Burden upon the poor Hen. Then is she oblig'd to perform several painful Employments, she must Sit upon her new hatch'd Little Ones and she must feed them. Sometimes she is restless and calling to the Cock, who often carries her nothing; sometimes she is in a Passion, rushing out of her Nest, either to beat the Cock, who refuses to supply her with Food, as she requires, or to fetch it herself to the poor Little Ones; like the Bee, which in a curious Plat of Flowers seeks the moisture that is upon them, and loads herself with it to carry to her Hive.

In the same manner you may see the Hen search all Corners of her Habitation, to find the most juicy of all those sorts of Food allow'd her, to carry it to her Young. She may be said scarce to regard herself; she only lives and acts for those Little Ones, always fearing she has not given them enough, and so she spends and wastes herself for them; insomuch, that sometimes you may see her decline and fall so Sick, that she either dyes, or at least recovers with much Difficulty. I am apt to flatter my self, that the Reader, having had this short view of all the Trouble and Fatigue a Hen endures upon this occasion,

occasion,, will readily grant that this suffering far exceeds the others of Laying and Sitting.

Those therefore, who desire to spare one Hen more than the rest, either in regard that she is tender, or that she is more beautiful and higher priz'd than any others they have, must manage after this manner. When you will make this Hen Sit, putting her into a Hut with a Cock, you must give her a Nest ready made, with something more to add to it, to the end that if she does not like she may alter it. When she has lay'd all the Eggs for the first time, they must be put under her to Sit on them seven days, and then look in to them to see whether they are Addle; if so, throw them away; but if good, put them under another Hen to Hatch. Let the first Hen rest two days; after which give her another Nest, made like the first, and when she has sat 5 or 6 days on her second Parcel of Eggs, take them away, and give her others that are ready to Hatch. Let her rear those Little Ones, which are none of her own 12 days, if she be a good Feeder; for if she is not, the Eggs ought to be taken away the day before they are to be Hatch'd. When you have taken away the Little Ones, to bring them up by Hand, let her rest two days, and then give her the third Basket, with the Nest

Nest ready made. Put a little very fine Hay into her Hut, to finish her Nest to her own Fancy. When she has fate upon her Eggs 12 days, take away and give them to another to Hatch, and put your Cock and Hen into a little Cage together, and leave them there till they begin to Moulte, when you may part them without any Danger. By this means you may perceive your Hen is not fatigu'd with thrice Sitting, and thus she will live long and have Strength to overcome the Distemper of Moulting, which carries off most of them, as being before too much weakned.

CHAP. XVII.

Of the several Distempers incident to Canary-Birds.

C*anary-Birds* during their Life time are subject to several Distempers, I know there are some of so good and happy a Constitution, that they may be said to pass the whole Course of their Life, tho' very long, without any Disease, except their Moulting, and very often they Sing during that very time, as loud, and as long,

long, as the rest of the Year; but being also well assur'd, that there are very few of that strong Constitution, I have thought fit here to set down the Distempers all *Canary-Birds* in generally may be subject to. I will begin with their most common Diseases, and conclude with those that are not so frequent.

Canary-Birds being thus liable to many Distempers, their Diversity causes various Effects, and produces several outward Signs or Indications, which if conceal'd, the Occasion of their Illness is unknown; so that for the most part no Remedy can be apply'd to them, being ignorant of the Malady, and of what is good or ill for it. The outward Signs or Indications, are therefore Circumstances absolutely necessary for knowing of the inward Distempers, which *Canary-Birds* as well as other Creatures are subject to.

In this Chapter I will speak of the understanding of the several Diseases that seize the *Canary-Birds*, and in the next I will prescribe the Remedies to be apply'd for the Cure of those Distempers.

The first Distemper I have observ'd in *Canary-Birds* is a Surfeit, a Disease the more dangerous, in respect that the Remedies apply'd have very often no other effect than the easing of the Bird, and prolonging its Life, without working a
perfect

perfect Cure. This Malady they often fall into a Month or 6 Weeks after they are Hatch'd. The Indication of this Disease is external; for you will see the *Canary-Bird* that has it very lean, his Belly transparent, big, very hard, full of little red Veins, all their little Bowels being generally sunk down to the extreme part of their Body. Some of them continue to eat well, notwithstanding this Disease; but they must infallibly dye very soon, unless proper Remedies be timely apply'd to this Disease. There are several Causes which may contribute towards this Disease, I am acquainted with two of the chiefest. The first is their Body being parch'd up within, by giving them too nourishing Food, whilst they were brought up by Hand, as for Instance, putting too much Sugar, or Bisket into their common Paste, as is still practis'd by several Persons, who kill them with Kindness.

The second is, that they relish so well all that is given them, when they begin to Feed alone, and are so well pleas'd that they can live without help, that they eat a great Quantity, and ravenously, whatsoever they have in their Cages; so that they generally Surfeit. When therefore you have any Young *Canary-Birds*, that are continually Feeding, to prevent this Distemper, take out of their Cages that
which

which you see them eat most greedily, giving them some of it now and then, without letting them be continually us'd to it. If nevertheless they fall Sick of this Distemper, you may make use of several things successively, which shall be set down hereafter.

Moulting is a Distemper no less Dangerous to *Canary-Birds*, than that we have last spoken of; there would be many more curious Persons in this Point, had not they quitted the innocent Pleasure of breeding such fine Birds, on Account of Disgust and a sort of Despair, seeing very often, in less than a Month, all the Birds dye, which they had been at the Trouble of rearing the best part of the Year. This Distemper carries off as many *Canary-Birds* as breeding of Teeth does Infants, great Numbers of which daily dye, for want of Strength to endure those sharp Pains; but as all Years are not equally fatal, and some Years scarce any *Canary-Birds* dye of Moulting, as for instance, when Autumn is fair and temperate: This ought to be an Encouragement to such as design to take this innocent Diversion of breeding them. Besides, if no *Canary-Birds* dy'd of Moulting, which is the Distemper that destroys the most, they would in a short time be more common than Linnets, and then their Numbers would in some measure

measure render them contemptible and of no Value.

I must confess I know no Remedy that can save them all from that Distemper; had I any such to my self, it would be a Treasure to me, and I would desire no other Employment for my Pleasure, and to get Money by.

At the Moulting time, which begins 5 or 6 Weeks after they are Hatch'd, and holds them above two Months, you will see them all huffed, melancholy, and often sleeping in the Day time with their Heads under their Wings; you will find the Cage they are in cover'd with small Down. The Young Ones cast their Down only the first Year and the second their great Feathers, as those of their Wings and Tails. At that time they are very uneasy; they eat but little and very often will not touch that which they like best when in Health. This is the dismallest Condition they are ever in; they are quite stripp'd of their Feathers, when the Weather is cold, as happens to the *Canary Birds* that come of the last Sittings, which do not Moulte till Autumn is well advanc'd, and sometimes in Winter. The Remedies to be apply'd to them shall be prescrib'd hereafter.

They sometimes grow very Sick of a Pimple growing on their Rump, commonly

Canary-Birds.

95

ly call'd the *Pip on the Tail*; in this case Nature must be left to work as much as may be; that is, leave them to break it themselves; but when you observe they are much swollen and huffed, and you are sure it is not occasion'd by Moulting, you must look upon their Rumps, and if you find that Pimple, they must be speedily reliev'd, as shall be said hereafter. Some of them are sometimes so Sick with it, that they have not Strength to break it themselves; and if they are not reliev'd, they certainly Dye. This is often occasion'd by Melancholy, because of being in a dark Place, as also from their not being purg'd often enough.

They are subject to have some yellow Scabs upon their Heads, and sometimes about their Eyes; when this Distemper spreads, nothing must be apply'd to it; Time and cooling Food will carry all that off.

They grow Sick and Lean through the multitude of Insects that breed in their Feathers, which is discover'd by seeing them picking themselves every Moment. There are several things to be us'd for this purpose, which shall be spoken of in the following Chapter.

All the *Canary-Birds* in a new Hut, sometimes fall Sick and sometimes Dye soon after they have been put into it; those

those concern'd give them Forty several things to recover their Health, but all in vain; the Cause of their Distemper is inward, which is the reason why many of those who have been longest acquainted with them have not found it out. This is certainly occasion'd by the Huts being made of old Pipe Staves, which have long serv'd to hold Strong Wines; so that notwithstanding the Workmans having plain'd it, as well to take off the Tartar that sticks close, as to smooth it, still the Wood retains a strong Scent, which tho' it be not always smelt by others, is still the natural Cause of the Young *Canary-Birds* falling Sick, being dos'd and as it we drunk with that scent of Wine, and most of them Dye in a few days; for tho' the Old Ones may perhaps grow familiar with it, and live in that new Hut, yet the Young Ones being much tenderer than they, Dye almost as soon as Hatch'd, and we are apt without reason, to blame the Hen, for having suffer'd her Young Ones to Dye so, for want of Food.

The most natural Remedy to prevent such Accidents, is not to employ Workmen, who out of Covetousness make use of worse Wood than others, without considering what a considerable loss those Huts may occasion to such as make use of them. I know their Profit in it is con-

considerable, for an old Cask they buy for 6 or 7 Pence at most, affords them Stuff enough to make a Hut, which they frequently sell for Half a Piece. This is Temptation enough for them, not to regard what unlucky Accidents they may occasion to those who make use of such Huts.

You may know those Huts that are made of Pipe Staves, if you observe whether that Hut you are about is made of 20 or 24 Pieces join'd together; for the Top, the two Sides and the Drawing out Board are each of them made of 3 or 4 Pieces, this with the Troughs, the sliding Boards, &c. may amount to 24 Pieces, which being every where fastned together with little Pins, it often happens that tho' the Hut is but little us'd, it drops in Pieces, like the Houses Children build of Cards, which when any one Card is taken away, fall all together in a Moment. This is for the Reputation of many good Workmen, who use only good Stuff, and very often sell it no dearer than those who often practise the contrary.

If, notwithstanding what has been said, you will make use of Huts made of Old Pipe Staves, you must ask the Workman, whether they are newly made? which he ought to tell you, if he has never so little

F

Honesty;

Honesty; and if the Hut is New, you must expose it for some time in a very Airy Place, that it may by degrees lose that ill Scent it has contracted, and then put your *Canary-Birds* into it. I know one who would have, not long since, gone to Law with the Person that sold him his Hut; because all his Young *Canary-Birds* dy'd in less than two Days, fancying that the Hut had been poison'd; but having undeceiv'd him, by demonstrating that the Occasion of his *Canary-Birds* dying, was no other than what I have here mention'd, he desisted; only assuring me that he would never make use of such Huts again. However he would needs make an Experiment of what I told him, putting two healthy Grey *Canary-Birds* into that Hut, they two Days after began to huffle up their Feathers, and would certainly have Dy'd, had he not raken them out quickly. To obviate all such Accidents, it is best to have such Huts as have been describ'd in the 3^d Chapter.

CHAP. XVIII.

*Several extraordinary Medicines
for curing Distempers of Cana-
ry-Birds.*

IT would be to little purpose to know what Distempers *Canary-Birds* are subject to, unless we were acquainted with the Remedies which may give them ease.

I have therefore thought fit to insert here the most sovereign Remedies, in order at least to give them ease, if we cannot quite Cure them, especially under their 2 first Diseases I have spoken of in the last Chapter, which are very heavy upon them, and so very much weaken those poor Creatures, that the Remedies apply'd have not the Success which could be wish'd, and we are often oblig'd, after much Care and Trouble to let Nature work, and after all our endeavours, to loose that which our Medicines could not preserve.

Several Remedies may be us'd for curing of the *Canary-Birds* that have the first Distemper I have mention'd, which is the Surfeit.

When you have a *Canary-Bird* that is

Surfeited, which will appear, if blowing up the Feathers on his Belly, you perceive his Bowels black and twisted, with other tokens I have describ'd in the foregoing Chapter. Then take a bit of Allum about as big as a Pea, and put it into his Water to dissolve, and renew the same every day, which continue for 3 or 4 days. Many curious Persons have found good Effect of this easy Remedy.

Again, when you have a *Canary-Bird* in this Disease, put a bit of Iron, as a Nail, into his Water, and change the Water twice a Week, still leaving the Nail in it.

Some Persons at Night take away the Sick Birds drink, and in the Morning give him Salt Water; the Bird fails not at first to drink some Drops of it. When you have seen him drink several times of that Salt Water, take away that Salt Water, and give him his common Water again. This must be done 5 or 6 days successively, and if you do not find the Bird recovers, make the following Composition. Give him some Milk boil'd with an equal Quantity of the Crum of Bread, take away his common Seeds, and give him some boil'd *Canary Seed*, in a little Pot into the middle of his Cage. Do this for four or five Mornings successively, and in the Afternoon give him his common Seed again in his Trough. At the end
of

of the 5 days, at fix of the Clock in the Morning, put about the bigness of half a Lentil of Treacle into his Water, and leave it till you have seen him drink of it once or twice. Continue this drink two or three days successively, after which prepare for him the following Food; Take a pinch of Millet, the same Quantity of Canary Seed, a little Rape Seed, with some Grains of Hemp Seed, mix them all together; then let these Seeds just be boil'd up, and throw off that Water to rinse them in cold; then boil a new lay'd Egg hard, bruise the White and the Yolk together, putting but a Quarter of it at most to the Seeds; add to all this a little bit of hard Bisket, a little Shell full of Lettuce Seed, with the same quantity of Pink Seeds; make all these into a Paste and give it to the Sick Bird. Give him also some Leaves of very yellow Endive, and continue this excellent Composition as long as your Bird is Sick. I know this Paste is troublesome to make it every day; but when a Man has a fine *Canary-Bird* he is fond of, the Trouble is not regarded, especially when he perceives that the Medicines apply'd do him good, and often quite recover him.

When a *Canary-Bird* is Surfeited, and you see his Bowels through his Skin, you must give him some pounded Walnut,

with boil'd *Canary Seed*, and after it a white Cabbage Leaf and some Sallery.

For a *Canary-Bird* that is Sick with Moulting.

He must be set in the Sun, and if the Sun does not Shine, put him into a hot Place, where no Wind can come, for the least Cold at that time kills him. During the whole Moulting time give him what follows, *viz.* some Wild Tansey, or Silver-Wort Seed, mix'd with a little Pink-Seed, in a little Gally-Pot, in the midst of his Cage; the next Day give him a little dry Bisket and Simnel; give him likewise some steep'd in white Wine; if he eats it, that will do him much Good. Spurt on him with your Mouth, three times a Week, that is, every other Day, some White Wine, and then immediately set him in the Sun to dry, or else before the Fire. If you find him very Sick, make him every Day swallow three or four Drops of the same white Wine, sweetned with a little Sugar Candy, or common Sugar. Put into his Water a little fresh Liquorice well scrap'd; it gives the Water a Relish, and does not over heat them. If you do not perceive your Bird mends, then without any more to do, put into his Cage, besides what I have mention'd, all other sorts of Meat, as hard Egg, both the White and the Yolk, some Simnel, a little

little Lettice Seed, bruis'd Hempseed, Canary-seed, some of his own Seeds boil'd, and so forth. It is frequently observ'd, that Sick Persons, who were given over, recover their Health and Strength by eating something that pleases them, tho' it be contrary to their Distemper. The same must be practis'd with *Canary-Birds*, refusing them nothing they can eat, when they are dangerously Sick, and leaving Nature to do the rest.

When your *Canary Bird* is troubled with that little Pimple on his Rump call'd the *Pip*, and you observe he leaves off Singing, and grows very Sick, take him into your Hand, and with the Points of a fine small Pair of Scissars, neatly cut off the Head of that White Pimple, then squeeze out the matter gently with your Finger, and immediately lay on the Sore a little Grain of Salt dissolv'd in your Mouth, and that will certainly cure the Distemper. If you perceive your *Canary-Bird* is in much Pain, because the Salt smarts, you may an Hour after lay upon the Sore a little Sugar dissolv'd in your Mouth, which will correct the Sharpness of the Salt, and quite heal the Sore.

There are several harmless Remedies for easing the *Canary-Birds* of those little Insects, some call Lice, that disturb them. The first way to prevent, is keeping of

them clean; that is, changing their Sand very often, cleaning the Hut, or Cage they are in, twice or thrice a Week. They must also have Elder or Fig-tree Sticks left them all the Year about, bor'd at small distances with the Point of a small Needle, all the Pith being taken out, and the Bark clean taken off, that they may be very smooth. Those Sticks are to be scrap'd or well shaken twice a Week at least, to get out the Insects, if any happen to be in them. You may also at Night put into the Hut a Piece of Linnen Cloth well whitened with Lye; and if there be any Insects or Lice, you will see them in the Morning upon it. But some *Canary-Birds* are frighted to see any Linnen in their Huts, when they awake in the Morning, because the White strikes their Eyes and scares them, and therefore this latter Remedy is not always to be us'd.

Before you put your *Canary-Birds* into a Hut, if it be an Old One, you must boil some fair Water, and wash it very well with the same scalding, which will kill all Insects and destroy their Nits, which lye commonly hid in all the Corners of an old Hut. The same is to be done with Cages, when they are old; and thus your *Canary-Birds* will not be tormented with those Vermin, which are often very troublesome

blesome to them, for want of being kept clean, and serv'd as has been here pre-scrib'd.

All curious Persons, who have many *Canary-Birds*, must always have an Infirmary; for it is rare that among a great Number some one should not be Sick; which cannot recover unless they are parted from the others; for besides that it will have no rest, the several sorts of Food you would provide to cure it, would be soon devour'd by the other Birds in the same Hut, which have no need of them. To obviate these Inconveniences, the Sick *Canary-Birds* must be parted, putting them into a Cage made on purpose, and call'd the Infirmary. A *Canary-Bird* Infirmary is no other than a pretty large Cage, cover'd on the Top, at Bottom, and on two sides with a Piece of thick Red or Green Serge, so that no Light may come in but at the Front. The enclosure of this Infirmary is to be made of small Ozier and not of Wire, because this is always Cold and Damp. Set this Cage in the Sun, if it be Summer, and in Winter in some Place where there is Fire always. Take heed of exposing them where there is any Smoak, which is very hurtful and often kills those that are in perfect Health; for the Smoak getting into their Throat, especially when they Sing, stifles them in

a short time. A *Canary-Bird* put into such an Infirmary as I have here describ'd, is half Cur'd, and giving him besides, some small matter that is proper for his Distemper, you will soon see him recover his lost Vigour; for they require Warmth in all their Diseases, and it is plain that they cannot feel Cold in such a sort of Infirmary. The Troughs of this Cage must be plac'd in the Bottom, after the manner of those I have describ'd in the Chapter of Cages and Huts. Then give them all the Medicines proper for the Distemper you think they are afflicted with, which have been here set down at large; and if notwithstanding all the aforesaid Precautions, you have any *Canary Bird* that loses his Natural Heat, which you may perceive by his being Melancholy, and sleeping continually without Feeding, and Sitting in the middle of the Cage, with his Beak under his Wing, then take him immediately, and having made him swallow two or three Drops of good white Wine sweetned with Sugar, which is not to be administer'd unless in cases of Extremity, put him alone into a little Cage that has a Piece of fine Lambs Skin at the Bottom and about it, and so let him rest all the Night, hanging the Cage in some warm place, as at the Head of the Bed you lye in. The next Morning take out
the

the Sick Bird, and put him into a little Cage, close cover'd, without any Perch. I have in a manner rais'd some from Death to Life by this Method, which is neither Difficult nor Troublefom, especially to one that is fond of his Birds, and the more when he perceives that his little care is not lost, finding very often a *Canary-Bird* to live several Years, which others, less skilful than himself, would have given over as certainly lost. Do not put your Sick *Canary-Birds* among the others again till they are perfectly recover'd.

All Creatures having their several ways of cleansing or purging, I believe it will not be thought amiss, that I should here say something particular of the purging of *Canary-Birds*.

Purging of *Canary-Birds* consists only in changing for a day or two their ordinary Food, which is Rape Seed, Millet, *Canary Seed*, Hemp Seed, &c. and instead of all these giving them only Rape Seed, Lettice Leaves, Chickweed and Groundfel. You may also give them some little Raddish Leaves, as also Beet Leaves; but when the Season for all those cooling Herbs is over, give them in lieu thereof good Melon Seeds shell'd and Lettice Seed.

There are two ways to know when your *Canary-Birds* want Purging.

The first is, when they Dung with much difficulty,

difficulty, which is a sign that they are very hot within.

The other is, when you perceive that they are continually throwing out the Seeds you put into their Troughs, which shews that they Eat very little.

These two tokens, without losing time to mention many more, may suffice to convince you that your *Canary-Birds* are in much need of being Purg'd. During the two days when you give them the things above mention'd, you must also give them a little Sugar Candy, or common Sugar in their Water. These things may be given them twice a Month, and this Precaution will prevent their being full of Humours, they will always Sing Merrily, and have a good Stomach.

CHAP. XIX.

Other Distempers Canary-Birds are subject to, with the Cures.

C*anary-Birds* are subject to some Diseases, in which they may be help'd without any Trouble. For instance, when they are out of Order by being too well Fed

Fed and growing too Fat, you must then take away all the sorts of Nourishing Food usually given them, as Canary-feed, Millet, Hempseed, Simnel, Bisket, &c. and in lieu thereof give them nothing but bare Rape-feed, and if you perceive they can scarce eat it, to encourage them you must steep it for some Hours, and having thrown off the Water you may give it them; for the Shell being then somewhat softned they will eat it very well. You may continue this Diet till you perceive they are somewhat brought down.

They are subject to have some yellow Scurf or Scabs on their Heads; when it is not spread, nor thicker than a Hempseed, you may open the Sore with the Point of a Pair of Scissars, making a little Incision to let out the Malignity, which sometimes rises like an Excreescency, and then soften it with something that is moist, as Oil of sweet Almonds, sweet Lard, Capon's Grease, or fresh Butter; but when the Evil is spread all over the Body, make use of the Medicine I have set down in the 17th Chapter.

Canary-Birds sometimes suffer much, for want of a little Precaution.

Sometimes a *Canary-Bird* is put quite out of Order by being handled too roughly. You will hear him when he is in your Hand make a Noise like the snapping of a Finger.

a Finger. This Cry, or Noise is follow'd by some Drop of Blood coming from his Beak, and then you may see the Bird as it were stunn'd and not able to extend his Wings. You must put him again immediately into his Cage, cover him with some thin transparent Cloth, and hang him in some Place out of any Noise of People, that he may not be disturb'd; set his Meat and Drink on the Bottom of his Cage, taking away the Perches, and give him good Food; if he survives 24 Hours, there is Hope that he will not Dye, of it, but will only be gaunt and thin. This generally only happens to *Canary-Birds* that are Wild and Unruly, being commonly those that have been rear'd by both Cock and Hen. To prevent this Accident you must use Precaution, drawing near the Cage, out of which you would take the Bird, by Degrees; you must give him Notice at a distance, either with your Mouth or Hand; for if he be taken by Surprize, tho' he does not fall into the dangerous Distemper I have spoken of, it often happens that when you miss taking him at First, he flutters about the Cage, till he breaks his Head or Wings. When you would take a *Canary-Bird* that is in a Volery, you may use a sort of little Net made purposely to catch them. Some make a little sort of Trap Cage, which

Canary-Birds.

111

which they set into the Volery, with some sweet eatables in it, as Simnels, or Biskets; the *Canary-Birds* soon run into it, one after another, and sometimes several of them together. Those that are taken in it you put into a Cage, and return the Trap to the Volery, till such time as the Bird you would Catch is in it, and then you return all the others you have no Occasion for into the Volery. I find this a very convenient way of taking *Canary-Birds* out of a Volery; and must own this Method does not scare them, and consequently they are out of Danger of all those Accidents above mention'd, which happen when taken by Hand; but I am sensible of an Accident which may happen, and therefore discourages me from practising this way, and that is, that it may fall out, that as the Trap is falling to shut in the Birds that are within, others may be walking upon and about the Trap Door, not being able to get in till the many others that are before within are clear, and the Trap Door falling in the mean time very quick, before the Birds that are on the Edge of it can be gone, they are in great Danger of having their Claws broken, and sometimes of being kill'd. I leave the Curious to make Choice of what they shall think most for their Conveniency,
and

and least hazardous, for killing their *Canary-Birds*, when they are to take any out of their Voleriers.

There is still another sort of Distemper incident to *Canary-Birds*, which may be call'd a Drooping, because the *Canary-Bird* that has it, is puff'd up and swell'd; his Body appears full of little red Veins, his Stomach clings to, he Eats little all Day, and does nothing but throw about the Meat with his Beak. This Drooping may proceed from his being in a Dark Melancholy Place, or by several Cocks being in the same Cage, and taking such an Aversion to each other, that it makes them fall into this Malady. The Remedy is, parting of them into several Cages, if you suppose the Distemper proceeds from that Cause; if it be from being in too Dark a Place, you must mend it, by putting them into another that is more lightsome, to make them Merry and recover their Health. Till they are quite recover'd of this Disease, their Diet must be mended, giving them some little Sweet to eat, and putting some Liquorice into their Water.

It is a great Mistake to say, that *Canary-Birds* have the Distemper call'd the *Common Pip*. That which is commonly taken for the *Pip* is a sort of Canker that grows on their Beak, and proceeds from
a great

a great Heat in their Bowels; they may be cur'd in a few Days, by giving them several sorts of things to cool them, as Lettice-feed, and put a Pinch of Melon-Seeds into their Water for three or four Days; and when you perceive they are a little better, take away that Water, and give them fresh, with a little Sugar-Candy in it, and continue the same for five or six Days longer.

Canary-Birds are also subject to a Looseness, which is easily perceiv'd when their Dung is more liquid than ordinary. They are then continually shaking and clapping to their Tails, and themselves languishing. When you observe this, you must pull their Tail Feathers, and those about their Fundament, and anoint that Part with Oil of sweet Almonds, or fresh Butter, and then give them Lettice-feed and Melon Seed shell'd, for four or five Days; as also the Yolk of an Egg boil'd hard, leaving them very little of their common Food, especially the first three Days.

It often happens that *Canary-Birds* break their Wings, or their Legs; when either of these Accidents happen you must use the same Medicines. You must in either case put your Bird into a little Cage with some Moss, or fine Hay; take away his Perches, setting his Meat and Drink on the bottom of his Cage in a corner. Having

ving no Perch, he will not hurt himself by striving to get up. You must not bind their Feet, tho' they be broken, because it would cause an Inflammation in the Ligature. Hang him up cover'd in some By-place, for fear lest he happen to make his broken Foot, or Wing, worse, when any Body comes near his Cage; and the broken Foot being thus left to itself, Nature, which is the best Physitian, will soon heal it.

The Falling-Sickness is very dangerous to *Canary-Birds*, but they seldom are liable to it. Of all Birds the Gold-Finches have it most. When this Distemper happens to seize your *Canary-Birds*, if they escape the first time, you must pare their Claws and sprinkle them at least twice a Week, with strong Wine lukewarm; you must not let them Sit to Hatch, nor teach them with a Flageolet, but set them often in the Sun, to chear and make them Merry.

Sometimes *Canary-Birds* fall Sick by being too much heated inwardly; you must then take away their white Seeds, as *Canary Seed* and *Millet*, as also *Hemp Seed*, and for a Fortnight give them nothing but *Rape Seed*, *Lettice Seed*, *Chickweed* and *Groundsel*, if it be the proper Season, as in *May*, when it begins to be thorough ripe. You may also give them
some

some Raddish Seeds and other cooling Herbs. That which at one time is good and Physical for them, is Poison at another. Some Persons, as soon as they find any Chickweed, or Groundsel in Gardens, even before the Spring, immediately bring it to their *Canary-Birds*, who feed on it freely, and often dye in a few days after. That is then the only cause of their Death; the Chickweed, or Groundsel you so eagerly gather for them, is perfect Poison to them; because that Chickweed, or Groundsel, is in a manner Wild, being produc'd only by the Snow or Water, and not ripe; the Sun which has not yet shin'd on it, has not yet taken off its Crudity, and of Consequence it is rank, for which reason it does not Digest in the Stomachs of the *Canary-Birds* you give it to, and therefore they sometimes Dye in a few Hours, without your imagining the cause. Tho' you should think those Greens very ripe, and you had seen the Sun shine on them several days, yet you ought to give them to your Birds very sparingly, especially the first days; because those Birds having been the fix Winter Months without eating any, it would disorder their Bodies, being a change of Diet, which will certainly Gripe them, and occasion such a Distemper as is often very hard to be cur'd. I know one, who after he had
pleas'd

pleas'd himself all the Winter with the Thoughts of having several Couples of fine *Canary-Birds* to Lay and Sit in the Spring, lost them all in less than a Fortnight, in *March*, by giving them Greens, which a Peasant had brought him out of the Country. By this you may judge how dangerous it is to give *Canary-Birds* any thing out of Season.

When a *Canary-Bird* has an Asthma, you must give him some Plantain Seed and hard Bisket, steep'd in good white Wine. This Distemper is known when *Canary-Birds* very often in a day make a Cry, which they fetch up from their Stomach.

Canary-Birds are subject to another Disease, which some curious Persons call, *the cracking of the Skin*, and then their Voices quite fail them. This commonly befalls them after Moulting, when they have been three Months without Singing. They sometimes lose their Voice so far, as either to Sing very low, or not at all. You must then give them good things, which may moisten their Breasts, as hard Yolk of Egg cut small, with some soft Crums of Bread. You must put into their Water some fresh Liquorice well scrap'd; which gives the Water a Relish, and moistens their Throats.

When a Hen that has Young Ones happens to Sweat, which appears by the Feathers

thers on her Stomach and Belly being wet, as I have observ'd before, the Young Ones then under her, are in danger of being stifled, and you may see that their little Down does not come out. Curious Persons use several Remedies for this Disease, some of which I will here set down.

Some put a little Salt into half a Glass of fair Water, and when it is dissolv'd, they take the Sick Hen out of her Nest, and wash all her Belly with that Salt Water, and having so done for about half a Quarter of an Hour, they dip the Hen in fair Water, to take off all the Salt, and then putting her into a little Cage in the Sun, or before the Fire, she prunes herself and grows dry in a Minute, and then she is put again into her Hut.

For my part, I make use of the Bone of that they call the Cuttle Fish, which Jewellers also make use of, which I reduce into fine Poudre, and with it rub the Belly of the Sweating Hen. I find that dries her and takes off the gross sweat. This must be done every three Hours, till the Little Ones are 5 or 6 days old. If this seems to be too troublesom, you must do what I observ'd in the 13th Chapter.

These are the Distempers *Canary-Birds* are most subject to. There are some more, which I shall not mention, because it is very rare for them to be Cur'd, as
when

when they grow Blind with Age, or Gouty for the same Reason. There is no other Cure but keeping of them warm, till they have paid their last Debt to Nature.

CHAP. XX.

What sorts of Birds may be coupled with Canary-Birds, and of the Mongrels they breed

IT being natural for Man never to be Satisfy'd with what he has, but to despise what is in his Power, and ardently to desire whatsoever is out of his Power; curious Persons at present act accordingly in respect to *Canary-Birds*. They are not satisfy'd with having Abundance of *Canary-Birds* of the most beautiful sorts; but are for altering their Nature, and most of them take pains to make *Canary-Birds* couple with Birds of another sort, whose Young are call'd Mongrels; but where one succeeds, there are very many who make nothing of it, for want of knowing how to Couple their Birds right. Here follows what is to be observ'd, if you would

would succeed in breeding of Mongrels.
What Birds may be coupled with Canary-Birds.

Most of the Birds that Warble and Sing, as Yellow-Hammers, Chaffinches, Linnets, Gold-Finches, and many more, which would be too tedious to mention in this place, may be coupled with *Canary-Birds*, but it is requisite to be inform'd of many particulars, in order to succeed in having Mongrels. It is requisite to this purpose, that the Bird design'd to be Coupled with a *Canary-Bird*, have been brought up by Hand, and be us'd to no other Food than what is generally given to *Canary-Birds*, to the end that when they are shut up together there be but one sort of Provision made for them. For example, a Gold-finch is us'd generally alone to Eat nothing but Hempseed, such a Gold-finch so bred, cannot be coupled with *Hen Canary-Birds*, without great Hazard that one of them will Dye, for if you deprive the Gold-finch of his Hempseed, when he is in the Hut with a *Hen Canary-Bird*, and give him only Rape-seed and Millet, that Change of Diet will make him Sick, and may happen to kill him. If on the other Hand you leave him his Hempseed, as before, the *Hen Canary-Bird* that is with him, will eat so much of it as will make her Sick, because the Hempseed being

too hot a Food for her, must of Necessity be very hurtful to her. The same I have said of the Gold-Finch is to be understood of all other Birds design'd to be coupled with *Canary-Birds*.

Moreover, the Bird you put up with your *Canary-Bird* must be at least 2 Years old, especially the Hens, which scarce ever Lay the first Year; therefore many Persons being ignorant as to this part, have been quite discourag'd, having several Years coupled other sorts of Birds with their *Canary-Birds*, and never had any Young Mongrels.

It is also requisite to put those Birds you would have Sit, several Months before into the same Volery with your *Canary-Birds*, that they may grow acquainted by degrees, seeing one another long before they come to Couple.

Those Birds must be made as Tame and Familiar as the *Canary-Birds*, by keeping them in a low place, where People are always, that they may not be Wild, when you must come to supply them with what they want.

The general Method is to put a Hen *Canary-Bird* and a Cock Gold-finch, Linnet, or other; but for my part I am for the contrary, that is, a Cock *Canary-Bird* and a Hen Gold-finch, Linnet, or other, because, for the most part the Male com-
municates

municates more of his Kind to the Breed than the Female; and consequently the Mongrels that come from a Cock *Canary-Bird* will be more Beautiful, and Sing better, than if they came from a Hen *Canary-Bird*.

As for the Mongrels that come from these Birds, they are not all equally Beautiful, and some of them are but very indifferent as to their Feathers and Singing.

For instance, the Mongrels from a Yellow-hammer, are a little Blewish, and the Cocks of them Sing very ill, especially when the Sire is a Yellow-hammer, and the Dam a *Canary-Bird*.

The Cocks which come from Linnet Mongrels Sing much better than any others, but that is all, for their Plumage is very ordinary.

The Green Bird tho' it warbles, will bring no Mongrels, his loud Note and gaping Throat, when in love, frights the *Canary-Birds*, and they run away as far as they can; so that it is rare to see them couple with *Canary-Birds*, unless it be an old very sprightly *Canary Bird*, that has been Bred among those Birds; and there may be some hopes of having little Mongrels.

If you would have beautiful Mongrels
G and

and good Singers, they must be bred out of a Gold-Finch, which is the beautifullest of Birds as to his Plumage. They may be said to be no less charming to the Ears, than they are agreeable to the Eyes; the great Plenty there is of them occasions their not being valu'd as they deserve. Those that build their Nests among Thistles are distinguish'd above the rest, as being stronger, sprightlyer and better for Singing than others. They differ something from the rest in the Colour of their Feathers, which are somewhat darker than those that are bred in other places.

If you would be successful in breeding such fine Mongrels, you must couple them thus: Take a white Cock two Years old, and that has not been coupled with Hen *Canary Birds*, for most *Canary-Birds* are not like our Nice Persons, they do not love Change; and seeing a Hen whose Notes and Colour are strange to him, he might happen to spend the Season without Treading that Hen. Besides that Hen Goldfinch must have been bred up by Hand, or at least it must be very long since she was taken, that she may be grown Tame and Familiar, and us'd by degrees to feed on Rape Seed and Millet, with very little Hemp Seed, for the reasons afore laid down.

You must not fail from time to time

to put some Thistle Seed into the Huts where Gold-Finches are Sitting; for they are very fond of that Seed, which is in a manner their first Food.

Put this Mongrel Pair together into a little Cage a Month sooner than the others, that they may have time to be acquainted before they come to make love.

Thus you see you may expect a fine Mongrel breed; for the Cock *Canary-Bird*, whose kind commonly prevails, will produce much White, and the Hen Gold-Finch communicating a little of her several Colours, the Young Ones will be worth any Money. I do not say the contrary may not be done, that is, to put a Cock Gold-Finch to a White or Copple Crown Hen *Canary-Bird*; but the kind of the Cock for the most part prevailing in the Breed, as has been said, the Little Ones bred by this latter Couple are almost all Gold-Finches. The Young Ones that come from these mix'd Birds, often breed others the next Year, contrary to the Opinion of him that has writ the contrary; and this second Mongrel Race are so beautiful that they cannot be too high rated. Such are those I saw the last Year in this City, which were sold for 500 Livres, that is the Mongrel Cock and Hen and three of their Young Ones. Nature had never

produc'd any thing so beautiful in Mongrel Birds. I will not here describe the Variety of Colours they were adorn'd with; I think the mentioning of the Price is sufficient for every one to guess at their Beauty.

All the little Cock Mongrels that come from these different sorts of Birds, are to be put under old *Canary*-Birds that are smart Singers, that they may instruct them in their Natural Notes. The same is to be observ'd as to Young *Canary*-Birds; that is, there must always be three or four Old *Canary*-Birds that Sing well, in the Volery with them, to teach those Young Ones, who will readily learn what they hear.

If you will bring up Young Linnets by Hand, and when they can feed themselves, put them to those Cock *Canary*-Birds I have just now spoken of, those Linnets in less than half a Year will Sing almost as well as your *Canary*-Birds, and will take the same Notes, so that if you did not see them, you would be apt to take them for true *Canary*-Birds.

There are several ways of knowing the Grey, Yellow and Ash *Canary*-Birds, &c. when they are of the Coppie-Crown breed.

1. By their having some white Feathers in the Tail.
2. By their having little white

white Spurs. 3. By their having Down, which you may know by taking the Bird in your Hand, and blowing gently on his Stomach and Belly, you will perceive a little white Down, which of consequence differs in Colour from his natural Plumage. Some *Canary-Birds* have much more of this Down than others; and therefore the Curious call the latter *Little Down Canary-Birds*, that is, that it appears thin; and the others are call'd *Great Down Canary-Birds*, that is, where there is much of it. This Down does not generally come till after they have Moulded. I must add, that there are some *Canary-Birds*, which are of the Copple Crown Breed, and yet have none of those marks I have above mention'd; so that you must rely upon the Honesty of the Person who Sells them you as such.

There are some *Canary-Birds* who have all the three marks I have spoken of, but one of them is sufficient, to perswade you that they are of the Copple-Crown Breed.

CHAP. XXI.

Why most of the Canary-Birds brought us from Germany, are apt to Dye soon after we have bought them.

Some *Germans* resort to this City twice a Year, that is, in Spring and Autumn, who bring over Thousands of *Canary-Birds* from *Tirol*, an Earldom in the Circle of *Austria*, and the Southern Part of *Germany*; for it is not to be believ'd, as some still do, that the *Canary-Birds* they bring are bred in the cold Parts of *Germany*. As soon as they come, People resort to them from all Parts; some out of Curiosity, to see whether they have any *Canary-Birds* that are not of the common sort; others go to lay out their Money. These last are better receiv'd than the former; but when you go to ask Questions of them concerning their *Canary-Birds*, or other such like Affairs, without Buying any thing, they give you a very bad Reception, and in short, they use you very roughly, making good the Proverb, *Point d' Argent, point de*

de Suiffes, or *no Money no Suiffes*; but as soon as you show them that precious Metal, without which the most Ingenuous Man is not valu'd, I say, when they perceive you come to Buy some of their *Canary-Birds*, they receive you very courteously, and in their broken Language express themselves very much your Humble Servants; and that they value your Money no less than your Person.

I find two Motives, which incline Persons newly addicted to the Curiosity, and not well acquainted with *Canary-Birds*, to Buy them of the *Swiss* or *Germans*.

On the other Hand, there are two Reasons which prevail with those who have been longer acquainted with *Canary-Birds*, not to Buy of them.

Not to spend much time, I must say in a few Words, that the first Motive young Dealers in *Canary-Birds*, have for buying them of the *Germans*, is only Covetousness; because it must be granted they sell Cheaper than the City Breed-ers.

The 2d is, that because they have not seen *Canary-Birds* bred here, and the others come from far, they imagine there is something more choice in their Notes or Plumage than in those bred here. These

Motives weigh enough with them to Buy many of those *Canary-Birds*; but in this particular they are like those the Scripture speaks of, where it says, they have sow'd much and reap'd little; for of a Dozen they Buy, they can scarce show two alive six Months after.

This is the first Reason which discourages Men who have long been acquainted with *Canary-Birds*, from following their Example; for say they, the *Canary-Birds* the *Germans* bring, changing their Climate and being Fatigu'd with coming so far, and many of them in little Cages, fall Sick soon after they arrive, and no Medicines can recover them; and this may be call'd the Country Disease; for had they been left in their own Country, they had not Dy'd, as they do here daily; for where one Lives six Dye.

The second Reason they alledge is, that those *Canary-Birds* have been bred up with other sorts of Food than what we give them. When their usual Food is chang'd, for want of knowing what is proper for them, they cannot for the most part be enur'd to it, and that it is that occasions their Death. The *Germans* do not care to tell what Food is Natural to them, because dealing ingenuously would not be to their Advantage, for they

they would sell the fewer the following Years, great Numbers surviving, which being coupled would Multiply so fast, that in a short Time there would be no Vent for their Birds, the City being Stor'd with them. However they were not altogether so Cunning, in bringing along with them the first Years so many Hens, and having been too Covetous of Selling all their Cocks and Hens at their first coming hither, they now make little or nothing of it; because abundance of Persons have apply'd themselves to make the Cocks and Hens they have sav'd, Sit; and many in that great Number having prov'd successful, they now Sell to others, and hence it will come to pass, that in a few Years finer *Canary Birds* being to be had in this City, and as reasonable as those the *Germans* bring, People will not wait their coming, as they now do. But, if notwithstanding all I have said, any Man will still Buy *Canary-Birds* of the *Germans*, he must use the following Precautions.

First, he must Buy them in Autumn; I am sensible that he must then run the Hazard of their Moulting; but then, those Birds which survive, will be stronger and better settled, in order to put them to Breed, than those that are Bought in the
G 5 Spring.

Spring. Having thus out-liv'd all the Winter, they will be proof against the Change of Air; and being us'd to our Climate, there will be Reason to expect some Breed from them, rather than from those in the Spring, which have scarce had time to rest after their Journey, before it is time to Couple them in order to Sit. If you happen to Buy any in the Spring, and desire to save them, do not make them Sit till the next Year.

The 2^d Precaution is, to cover the Cage into which you put the *Canary-Birds* you have Bought of the *Germans*, for at least a Fortnight. Put them into a very close Place, as is usually done with Birds taken in Nets, because they have been cover'd all the while they Travell'd. If you set them at once into the full Light, it may occasion many ill Accidents too long to mention here, and particularly, battering themselves, as being too wild.

You must also continue some time giving them nourishing Food, and such as will warm them, as Hempseed, *Canary-feed*, hard Egg cut small with some of the soft of Bread. Put also a little common Sugar into their Water; for the *Germans* during all their Journey, give them all sorts of hot Food, that they may
the

the better endure the Fatigue of Traveling; not valuing if most of them Dye soon after their Arrival, their Bowels being in a manner Burnt out. Thus you may plainly perceive, that most of them Dye by the ill effect the Change of Food has on their Constitution; for, if you give the *Canary-Birds* Bought of the *Germans*, nothing but Rape-feed, as many here do, they will pine away by degrees and Dye, without your understanding the Reason of it, and therefore they must be us'd by little and little, to the Food usually given them here.

To conclude, the last Precaution to be us'd is, not to go for some days after the *Germans* arrive, to buy their *Canary-Birds*; because the two first Weeks after their Arrival, great Numbers of them Dye daily, of which they make no mention. Let the Dealers in Birds, who Buy many and Sell them again the same day, or the next, without regarding how soon after they Dye, go Buy them those first days. It is more likely that they will live when they have rested here about three Weeks, than if you take them the first days after their Arrival.

CHAP. XXII.

How advantageous it is to Breed up Canary-Birds by Hand; and for what Reasons the young ones of the First and Second Sitzings are to be preferr'd before the last.

MAny who have long been acquainted with keeping of *Canary-Birds*, may as much as they please, cry up those which have been rear'd by the Cock and Hen; for my part I affirm, with many others, that those which are brought up by Hand, especially when they have been Fed and look'd after, as is set down in the 8th and 9th Chapters, far exceed the others in Strength and Vigour.

In the first place, it happens that *Canary-Birds* rear'd by the Cock and Hen decline, either because the said Cocks and Hens are Sick, and do but half feed their Little Ones, or else are not able to perform it, having 5 or 6 to feed at once; which is the reason that in so great a
Brood

Brood they often neglect some of them, which for want of sufficient Food, Languish and Dye in a few days.

2dly, The Cocks and Hens are much eas'd, and do not spend themselves so much, when the Little Ones are taken from them after 10 or 12 days, which makes them longer Liv'd than those that every time they Breed are left to rear their Little Ones quite out: In fine, the Young Ones brought up by Hand, are much more Familiar than any that are rear'd by the Cock and Hen.

It has also been observ'd, that fewer of those brought up by Hand Dye in Moulting than of the others.

Besides, it gains one Breed of each Couple of *Canary*-Birds; for four Sittings may be expected, without spending them, during the four Months they are in Huts; because they Sit on their Eggs 14 days, which with 12 more before the Little Ones are taken away, make 26; and adding four days to Lay again, the whole amounts to 30; and tho' it should extend to 35 days, they may still Sit four times in four Months and a half; and after all that, will still be in a Condition to do the same the next Year; whereas, when they are left to bring up their Little Ones quite out, when they come to the 3d Sitting

ting, it is requisite to take them out of the Hut, because they are beginning to Moul't; and they are so much spent, that very often the Hens are fit for nothing the next Year.

I must inform you by the by, that it is better to Couple *Canary-Birds* early in the Year, especially when you have the Rising Sun to set them in and a clear Air, than to put it off too long. I know two ill Accidents may attend their Coupling too soon. The first is, when any little Frost happens at the time when the Hen is ready to Lay; for her Pores being then all clos'd, it happens that she cannot Lay her Egg, and Dyes, unless speedily reliev'd.

The 2^d is the Little Ones being Hatch'd in cold Weather, and then they are in much danger of not thriving.

When these cross Accidents happen, and you have put your *Canary-Birds* into their Huts; you must keep all the Windows close shut, and keep a Fire in the Place where they are, as long as that Weather lasts, and if there is any likelihood of its Continuance, you may take your Birds out of the Huts for a few days, and put them asunder into little Cages. If any one of the Hens has Laid, the Eggs must not be given her to Sit on, till the Weather alters.

On

On the other Hand, if you delay Coupling your *Canary-Birds* too long, for fear of such ill Weather, which does not frequently happen. I say, if you defer putting of them together too long, you will be oblig'd to part your Pair of *Canary-Birds* before the third Sitting, because of the too great Heat of the Season, which makes them come to Moulting, and then there is no more to be expected from them that Year.

There being Reasons *pro and con* for all things in this World, every Man, according to what has been said, may take his Choice, of running the Hazards he shall think fit, of what may be expected upon these Occasions.

It is said, that the First and Second Breeds are the best; but the Reasons for the same have not been yet assign'd.

For my own part, I find two principal; the first is, that the first and second Broods are always better Fed than the last; because the old Cock and Hen not being as yet tir'd, spare no Pains to Feed their Little Ones; whereas at the 3^d and 4th Broods, being quite tir'd out with following the same Employment so long, they are quite spent, and very often take so little Care of their Young Ones, giving them but very little Food, that they fall Sick

Sick and Dye before they can be taken from them.

I may say, that those who bring up the little *Canary-Birds* by Hand, are much like the Cocks and Hens themselves; for they are full of Care to bring up and feed the first and second Broods in due Time and Season; but when they come to the 3d and 4th Broods, they resemble the *Canary-Birds*, and grow weary of doing the same thing continually; and refusing to be any longer ty'd to Hours, they feed the Little Ones sometimes too soon, and sometimes too late; and being so irregularly look'd after, the Little Birds contract an ill Habit of Body, and Dye of the first Distemper they fall into, which is commonly their Moulting.

The last reason there is for preferring the first Broods before the latter, is, that the *Canary-Birds* coming into the World early in the Year, that is, in *April* or *May*, they Moulte either in *July* or *August* at the latest, being the most favourable time for that Distemper, because the Weather being then very hot, all the Pores are open, and the Feathers of those little Creatures drop off, and others come out faster and with more ease than at any other time. On the other hand the Little Birds that are Hatch'd late in the Year, as are those
of

of the 3^d and of the last Broods, which come in *July* and *August*, cannot Moulte before the middle of *Autumn*; and sometimes the beginning of *Winter*, when the Pores are shut, and consequently they have much Difficulty in Moulting, and very often all the Remedies apply'd to them, to help Nature prove ineffectual; so that the Cold hindring their new Feathers from coming out, nothing is to be expected but Death.

Be not of the Opinion of those, who say, that several *Canary-Birds* are not to be put together into one Cage; for if you were almost every day to clean as many Cages as you have *Canary Birds*, the trouble would overballance the Pleasure. You may therefore put two Birds into a large Cage. If they be Hens, you may put in four, because they live more peaceably together than the Cocks. You may put eight into a large Hut and into a Volery in Proportion to its bigness. If there happens to be an unlucky Cock among them, that pulls the Feathers off the others, as sometimes happens, take him out from among them and put him into a Cage by himself; but they must not be separated on account of any little Accidental Scuffles that happen among them, when they go no farther.

CH A P. XXIII.

Of the common Prices Canary-Birds are now sold for at Paris.

THere are very many Persons of all Ages and Conditions in this City, who do not yet keep *Canary-Birds*, and yet would be willing to have One or Two, to Divert them after their usual Affairs are over, with their sweet Notes; but most of those Persons still remembring the Price that *Canary-Birds* bore ten Years ago, without enquiring what they are now worth, believe it is still the same. Thus the Ignorance of some, and the Covetousness of others, are the cause that many go no farther than to Wish for *Canary-Birds*, without attempting to Buy them; my Design is to undeceive those People, by writing this Chapter for their Sakes, wherein I will set down the common Prices *Canary-Birds* now bear. To observe some order, I will begin with the Prices of the most common *Canary-Birds*, and conclude with those that

Canary-Birds.

139

that are by curious Persons reckoned the most beautiful and dearest.

A Common Grey *Canary-Bird* 3 Livres, 10 Sols.

A rough footed, or white Tail'd *Canary-Bird* 5 Livres.

An Ash *Canary-Bird*, rough footed, or white Tail'd, 5 *l.* 10 *s.*

A Common Yellow *Canary-Bird*, 4 *l.*

A Yellow *Canary-Bird*, rough footed, or with a white Tail, 6 *l.*

A Common Mottled *Canary-Bird*, 4 *l.*

A Motled *Canary-Bird* of the Copple-Crown Breed, 5 *l.* 10 *s.*

A Common Buff Colour *Canary-Bird*, 4 *l.*

A Buff Colour *Canary-Bird* with a cast of Gold Colour, 5 *l.* 10 *s.*

A Common Copple-Crown *Canary-Bird*, 6 *l.*

A *Canary-Bird* with an Ash Colour Copple-Crown, 7 *l.*

A *Canary-Bird* with a Black Copple-Crown, 10 *l.*

A *Canary-Bird* with a regular black Copple-Crown, 20 *l.*

A Common Lemmon Colour *Canary-Bird*, 15 *l.*

A Lemmon Colour *Canary-Bird* with a black regular Copple-Crown, 25 *l.*

All white *Canary-Birds* with red Eyes are so little valu'd, at this time especially, when

when there is Plenty of others, that I have not thought fit to speak of them, according to their several degrees of Beauty, the finest of them not exceeding the Value of 4 *l*.

The Price of *Canary*-Birds here set down, may vary upon two several Occasions. The first is, when those same Birds are Bought some days after they are Hatch'd, as many do, to breed them up by Hand; then at least the third part of the Price is to be abated, in every sort of what is here set down. For instance, a *Canary*-Bird that costs 15 Livres, when he can Feed himself, is worth but 10 Livres, if taken when he is but 10 or 12 days old.

The Price in like manner advances one third beyond what has been set down, when a *Canary*-Bird is Bought after he is pass'd all the Danger of his first Moulting, as in *March*, the time when he is fit to be put into a Hut. So a *Canary*-Bird of 15 Livres Price in *October*, will cost 20 Livres in *March* the next Year, and the like of all others.

That nothing may be omitted, I must take Notice, that there are *Canary*-Birds which Sell above the Price here set down: Sometimes one or two black Feathers in the Tail of a Coppel-Crown, or any other particular marks like Stars, answering uniformly

formly about his Body, will raise him to double the Price of the others. I will not here set down the Price of that sort of *Canary-Bird*, which is not common, and whose Value generally depends on the Will of the Person that is to Sell him to one that wants no Money, and is eager to Buy him. We frequently see curious Persons, who are well to pass in the World, and make nothing to give 3 or 400 Livres for a Couple of *Canary-Birds* they fancy.

I will not be particular as to the Price of Mongrel *Canary Birds*: Some of them are not worth so much as the natural *Canary-Birds*; but there are also others, whom Nature has form'd so Beautiful and regular, as if they had been Drawn with a Pencil. The Price of these is considerable, and such being very rare, he who has them, often meets with curious Persons who give him whatsoever he asks for them. As for instance, three little Mongrel *Canary-Birds* were some time since Sold here to a Curious Stranger for 500 Livres.

Nor will I spend time upon the Value of *Canary-Birds* Eggs; because the Selling of them begins to grow out of Date, tho' it was much us'd some Years ago. Those who first began to breed *Canary-Birds* in this City, made what they would of them, for they Sold the very Eggs at considerable Rates.

Rates. Some have been Bought for 10 Crowns a Piece, and very often they were Addle, or the Little One dead in the Shell. Now Money is grown somewhat scarcer, and *Canary-Birds* more plentiful, there are no such extravagant curious Persons.

Besides, it is common to be Cheated in Buying of *Canary-Birds* Eggs, through the Dishonesty of the Seller, as hapned to a Friend of mine last Year, who Bought *Canary-Birds* Eggs of a Man, that through Ignorance had Coupled two Hens together, whose Eggs can never be good for any thing, as is easy for any one to judge, having no Cock with them. And that Man, who Values himself still upon having a nice Conscience, made no Scruple of keeping the Money he had receiv'd; tho' a few days after he was convinc'd of his Ignorance, in Coupling one Hen with another, instead of a Cock, and very well knew that he who had bought the Eggs of him, hazarded his Money, upon the Supposition that those Eggs came of a Hen that had been Coupled with a Cock, as they are usually Pair d.

There are also some Persons so Deceitful, as to make you believe, that the Eggs you design to Buy of them, and which they shew you in a Nest under a Hen Sitting on them, are newly lay'd under

der her, tho' they have been Sate on 7 or 8 days, and consequently unlikely ever to prove good. This perhaps makes you have a Mind to Buy them, especially considering the extraordinary Beauty of the 2 *Canary-Birds* they come from, and very often they are the Eggs of another Pair of very Common *Canary-Birds*. Then whatsoever you Bid, the Sharper lets you have them, pretending it is to oblige you. Thus imposing upon your Credulity, he perfectly Cheats you; and if you afterwards come to complain that his Eggs prov'd Naught, he will answer, it is Pity, and he is very Sorry for it; but that such an Accident may happen to any Man, and so will endeavour to satisfy you with fine Words.

I have thought my self oblig'd to take Notice hereof for two Reasons.

The first, to discourage several curious Persons, who are not well Vers'd in this Affair, from Buying Eggs after this manner at Random, their Money being for the most part thrown away; because there are every where People ready to impose upon their Credulity, and to treat them as has been observ'd above; and they being thus fore-warn'd, will be the more upon their Guard, being first convinc'd of the Probity of those from whom they would
Buy

Buy their Eggs; and consequently will not grow weary of the innocent Diversion of *Canary*-Birds, when they see all they undertake succeed.

2dly, Because curious Persons being thus warn'd not to Buy *Canary*-Birds Eggs of all Persons without Distinction, they will avoid being the innocent Cause of the Fault committed by those who burden their own Consciences by wronging them of their Money.

It may perhaps be urg'd, that the Prices here set down by me, are of no Use, because the scarcity of Money hinders the Sale, and occasions their going now at moderate Rates. To this I answer, that there are now so very many curious Persons in this City who Breed *Canary*-Birds, that when Money shall become more plentiful, the *Canary*-Birds will be so in like manner, and that among the many curious Persons who will have Young *Canary*-Birds, there will always be several who will Sell them at an easy Rate, to dispose of the more, and to continue the reasonable Price I have here set down.

I would advise the Curious, who have not had long Experience in this Affair, not to Buy Young *Canary*-Birds for Breeders; because it often happens that such Persons being unskilful in the Business, and the
Birds

Birds too Young, they make nothing of it the first Year.

For my own part, when I want a Pair of *Canary-Birds*, I find that a Cock three Years old, and a good Hen of two, fit me better than two *Canary-Birds* Hatch'd within the Year, whose tender Youth is subject to many unlucky Accidents, too long to be taken Notice of here.

It is also to be observ'd, that when you Buy a Copple Crown *Canary-Bird*, tho' he be never so well mark'd, his Marks will wear off every Year at Moulting time, so that in Process of time he will turn quite White, without any Spots or Marks at all. For this reason curious Persons should be cautious how they give a great Price for a *Canary-Bird* curiously mark'd, since that Beauty wears off at every Moulting, and consequently his Value declines. I know there are some that retain their Spots long, and increase the Number of them; but such *Canary-Birds* are very rare.

As for the Charge of keeping a *Canary-Bird* a Year, it is Cheaper buying in his whole Provision at once, than by small Measures; and thus every Bird that does not Breed will cost 20 Sols a Year, and if it Breeds, reck'ning the hard Eggs, Simnels, Biskets, &c. which must be allow'd them at that time, the Expence will

H

amount

amount to about 30 Sols a Year. I know there must be some Management in the Distribution of their Seeds and other Food, for otherwise they spoil much more than they eat. Thus you see it is Cheaper keeping of a good *Canary-Bird* a Year, than a pittiful Chaffinch.

CHAP. XXIV.

*The Names, Qualities, and Prices
of several sorts of Seeds com-
monly us'd to Feed Canary-
Birds.*

I Think it will not be improper here, in a few Words, to set down the Names and Qualities of all the Seeds curious Persons use in Feeding of their *Canary-Birds*; and for observing of some Method herein I will begin with those Seeds which are most necessary for them, and will afterwards proceed to those they may live without.

The Names of Seeds proper for Canary-Birds.

Rape seed.

Millet.

Millet.

Hempseed.

Canary-feed.

Pink-feed.

Lettice feed.

Silverweed, or Tanfy-feed.

Plantan-feed.

Rape-feed is a very small round Seed; coming from a Plant of the same Name. I name it first, because it is the most necessary Seed for Feeding our *Canary-Birds*; and they may live with this alone, tho' they have none of the others, as has been said before. There are several sorts of it; one of them is bigger and blacker than the true Rape-feed, and commonly kills all the *Canary-Birds* it is given to, by reason of its Bitterness and other ill Qualities. The best is that which is smaller than this bad sort I have last spoken of. It is not quite Black, but inclines a little to Purple, is very sweet, and has none of the Bitterness of the other sorts; it is call'd *French Rape-feed*, that is the best that can be given to Birds. It has a Nourishing and Cooling Quality at the same time; so that a Bird which lives altogether upon this Seed, is not apt to grow so Fat as those that eat much of the others above mention'd.

When it is too old, as of 3 or 4 Years, it generally has no more Scent than Dust, nor scarce any Taste, and therefore does not so well Nourish the Birds that eat of it. On the other Hand, when it is too new, tho' never so good, it Surfeits them. It must be at least 6 Months Old before you give it to your *Canary Birds*, and to prevent being Cheated, you must Lay in a Years Provision before *March*: So that if it happens to be new, it will be at least 8 or 9 Months Old, so that it cannot be prejudicial to your Birds.

To be provided with Seed for a Year, without being oblig'd to Buy again within the time, you need only take 9 or 10 Litrons, that is something more than so many Pints for every *Canary-Bird* you design to keep, and you will find that enough for the Year.

Millet is a sort of small white Grain, round, and at least twice as big as the Rape-feed; the whitest is the best. There is a yellow sort which is only good for Poultry. The best Millet is that of the Province of *Anjou*, on the River *Loire* in *France*. It is sweeter and pleasanter than the Rape seed. Its Qualities are to Nourish, to Warm and to Fatten considerably, and therefore curious Persons must take heed not to give their *Canary-*
ry-

ry-Birds too much of it; nay sometimes they must be made to Fast, for the Reasons mention'd in this Treatise. This Millet serves for several other uses, which is not to our Purpose to take Notice of.

The Hempseed is the Seed of the Plant of the same Name. It is twice as big as the Millet, and Grey. The best is the midling size, and of a bright Grey. Its Quality is to Nourish, Fatten and Heat much more than the Millet last spoken of; and therefore very little of it is to be given to *Canary-Birds*, unless it be in the sharp Winter Season, at which time it is good for them. The best sort has somewhat of the Taste of a small Nur, and therefore the *Canary-Birds* are very fond of it.

The Canary-feed is a yellowish Grain, not so thick as the Millet, but longer, pointed at both Ends. Its Quality is to Fatten and Warm *Canary-Birds*. It has almost the same Taste as the Millet. Many curious Persons never give their *Canary-Birds* any of it, pretending that it Burns their Bowels; but it can do them no harm, provided they are not continually us'd to it, and have only a little Pinch given them sometimes.

The best Pink-feed comes from *Stras-*
burg,

burg, as has been found by Experience, that which grows about *Paris* loosing its Taste. It Blossoms about *May* and *June*. Its Quality is astringent, and therefore it is given to *Canary-Birds* that are Loose; the Taste of it is sweetish. Take heed when you buy this Seed, that they do not instead of it give you Poppy-seed, which is very like it, and will infallibly kill your *Canary-Birds*; all the difference between them is, that the Poppy-seed is Blackish, and the other Grey.

Lettice Seed is well known, as well as the Lettice, it is flat, long, and of a bright Grey. Its Quality is cooling, wherefore it is now and then given to *Canary-Birds* to Purge them. The freshest is the best.

The Silverweed is a Plant whose Leaves are like those of Coriander, but somewhat more Oily, and the Stem is like that of Rue. It is call'd Silverweed because the Leaf is white, but the Seed is red and very small. In *Latin* it is called *Thalictrum*. Its Quality is to bind the *Canary-Birds* that can Eat it; which many of them will not. It is put to several uses by Women, of which I shall not speak here.

The Plantain Seed comes from a Plant of the same Name, which grows like an Ear of Wheat, is small and blackish

ish. Its Quality is Nourishing and Warming, but it is very seldom given to *Canary-Birds*.

The Price of each sort of Seed cannot be positively determin'd; because it alters every Year, according to the Plenty or Scarcity of it. What I shall set down here, is rather to give curious Persons some Notion of the Value of the several Seeds, without their depending on it, and absolutely to fix it.

This being a dear Year, the Price I shall set down will rather be in Favour of the Seller than of the Buyer; because there is more reason to expect it will fall than rise.

A Boisseau, which is about a Peck and a Half *English*, or 20 Pounds Weight, 45 Sols.

A Boisseau of Millet, 30 Sols.

A Boisseau of Hemp Seed, 22 Sols.

A Litron, being somewhat above a Pint *English*, 3 Sols 6 Deniers.

A Litron of Pink Seed, 12 Sols.

A Litron of Lettice Seed, 9 Sols.

A Litron of Silverweed Seed, 28 Sols.

A Litron of Plantan Seed, 15 Sols.

The three first sorts of Seeds I have nam'd, *viz.* Rape Seed, Millet and Hemp Seed, must be very well fann'd and cleans'd before they are us'd; because they are ge-

nerally very foul, and full of little Stones and Dust. It may perhaps be ask'd, why I speak of Boisseaus, or Pecks, in mentioning the 3 first, and only of Litrons or Pints to the other 5. To this I answer, that those 3 first sorts of Seeds, are absolutely necessary for them, especially the 2 first of them, so that they may Live very well without the others, which are not proper for them, otherwise than to recover their Stomach, when they have lost it, or to help them in some Distempers I have fully spoke of in this Treatise. Those three sorts of Seeds are to *Canary-Birds*, as Bread, Wine and Flesh are to Men; for a Man who has those three sorts of Sustenance, may live long and in Health, without the help of any others, which are call'd the Dainties of this Life, and which often rather impair than preserve Health, especially when they are become habitual. By what I have said you may perceive, that a Litron, or Pint of the last five sorts of Seeds will serve you as long as a Boisseau, or Peck of the three first, being given to the Birds, only in cases of absolute Necessity. There are also many curious Persons, who never make use of all the sorts of Seeds I have mention'd; which did not however divert me from setting them down, for the Satisfaction

isfaction of those who do, or may hereafter use them. There are still several other sorts of Seeds given to *Canary-Birds*; but in regard that they live no longer than others that have them not, I have not thought fit to put the Reader to the Charge, or Burden his Memory with them, since he may do very well without them.

Rape Seed, Millet, Hemp Seed and Canary Seed, are generally sold in the Common Seed Shops; every one must make his Price when he understands the Seed, for I have seen some above 10 Sols dearer in the Boisseau than others, tho' the same sorts of Seeds. But as for the Pink, Lettice, Silverweed and Plantan Seeds, they will be found better and cheaper among the Flower Seed Men.

CHAP. XXV.

The use to be made of Canary-Birds, and how long they may live, when well manag'd.

Several who had newly apply'd themselves to the Curiosity of keeping *Canary-Birds*, have taken taken a Dis-taste to it the very first Year; because those who Sold them the Old Ones, which they put to Breed, had made them expect to have 15 or 16 Little Ones from each Pair; and having thus unreasonably flatter'd themselves with the Hopes of so great a Number, they have been quite discourag'd, finding scarce a third Part of what was promis'd them. He who expects but a small Number of Little Ones from each Pair, is agreeably surpriz'd, when there are more than he expected; and on the contrary, he is quite dejected, who finds fewer than he promis'd himself. We may indeed expect 15 or 16 Eggs from each Pair of *Canary-Birds*; and sometimes they Lay more; but to avoid being disappointed,

appointed, we are not generally to think that above one third of them will come to good, as well because of the Addle Eggs there are in that Number, as because some of the Little Ones do not come to good.

Thus to have *Canary-Birds* Eggs is something of a Beginning; if they prove good they afford hopes; when Hatch'd there is some Pleasure; and if the Young Ones out-live their Moulting, that Crowns the Work. I know there are some good Kinds which bring Considerable Breeds; but to avoid mistaking our reckoning, we ought to be satisfy'd, if taking one with another we have six *Canary-Birds* from every Pair alive after the first Moulting. I say taking one with another, because among several Couples put up to Breed, some Multiply very much, to supply the defect of other Couples, which make nothing of it all the Year, or but little at best, as I have observ'd in the 15th Chapter.

There are two sorts of curious Persons. The first are those, who have very many Couples of *Canary-Birds*, which they regularly put together to Breed every Year, making a sort of Revenue of the Little Ones. These Persons use the old *Latin* Proverb, *Omne tulit pun-*
ctum.

Etum qui miscuit utile dulci. That is, they know how to make a Profit of their Pleasure. In short, besides the Pleasure it is to them, to have such Numerous Broods of fine Birds, they make a considerable Yearly Advantage by Selling so many.

There were some Persons, even of Note, who drove an open Trade with them, the first Years after they came to this City; I believe it was not so much for the Profit they made of them, as for the Reputation of being successful in that new sort of Pleasure. But since many have had the same Success as they, and by that means the *Canary-Birds* are become more plentiful, they have left off that small Trade, to make Room for very many, inferiour to them in Birth, who have apply'd themselves to it.

The second sort of curious Persons are those, whose small Number of *Canary-Birds* afford them no other Prospect, but the diverting of themselves some Moments of the Day, after having spent the greater part of it in more solid and profitable Affairs than the devoting of themselves to *Canary-Birds*. There is no approving of their Practice, who the more *Canary-Birds* they have,

have, the more they covet, and the Number whereof often causes them to lose the best Part of the Year, by reason of the continual care they must take of them. That ought to be left to those who Deal in them, whose Business it is to Breed and Sell all sorts of Birds. I would therefore advise any one, who only keeps them for his Pleasure, never to keep above 5 or 6 Pairs of *Canary-Birds*; viz. 3 Pairs of Copple-Crowns, and the others plain Greys, or Greys with white Tails, to be as it were Nurseries, in Case of Necessity, to those that come from the Copple-Crowns. The Increase will be sufficient to serve himself, and to oblige some Friends. The continual Care and Trouble, which for at least one third part of the Year, attends a great Number of Sitting *Canary-Birds*, far out-balances the Pleasure and Satisfaction they can afford, especially if a Man has no other on whom he can repose that Trust.

I do not, however blame those, who every Year Sell a small Number of their Little Ones among their Acquaintance, to avoid increasing the Number of Pairs they design to make Sit the next Year for their Diversion; the little Money they

they make of them, only serving to defray the Charge of the Birds they keep for their own Satisfaction. It would be wrong, for any ignorant or malicious Person, to call them Traders in *Canary-Birds*, since it plainly appears by the Posts or Places they are in, that their *Canary-Birds* are not their constant Employment, nor afford them any settled Income, but only an agreeable Pastime; and that the little Money they make, by Selling, from time to time, what they have to spare, only serves to maintain their Volery; and as we cannot call a Citizen a Wine Merchant or Vintner, for Selling the overplus of Wine of his own Growth, after serving his Family, since we daily see many Persons of Condition, sell theirs, either by Wholesale or Retail, without being look'd upon as Vintners, as having other Places or Employments, which distinguish and raise them above that Employment; in like manner our curious Persons, may as long as they please, Sell what *Canary-Birds* they have to spare, after the Example of several Persons of Distinction, either on Account of their having some extraordinary Beautiful, or to employ the Money they get to Defray the Charge of their innocent
Diversions,

Diversion, without being liable to be call'd Dealers in Birds.

Only those ought to be call'd Dealers in Birds, who keep Publick Shops, and who are allow'd as Masters in their Profession, and value themselves upon it; or those, who having no other Income but what their *Canary-Birds* afford them, admit of all People indifferently to their Chambers, to sell them their Birds for Better or Worse.

He who is resolv'd to venture upon having a Brood of *Canary-Birds*, if he will be sure to succeed, must have more than one Pair; for if the Cock or the Hen of that Pair should happen to fall Sick, you are quite spoil'd, especially when they have either Eggs, or Little Ones, as not knowing what to do to preserve the Brood; but when you have several Pairs, there is commonly some Hen that has been Sitting or Feeding much about the same time, and to her you give the Eggs, or the Little Ones of the other that is Sick, and by that means you save that which must have been lost if you had but one Pair of *Canary-Birds*. When you have any Cock, or Hen, over and above what you put up to Breed, be sure to put it into another Room distinct from the rest,

rest, so that they may not hear them Sing, otherwise it may happen, that your *Canary Bird* hearing the others that are in their Huts, may grow hot in love, and will Dye of that Distemper in a few days, for want of being Pair'd.

If you would proceed orderly, without burdening your Memory, which ought to be better employ'd than on the Thoughts of *Canary-Birds*, you must every Year make a little white Paper Book, to set down the precise day you gave the Hens their Eggs to Sit on; the day they are to Hatch, if good; I speak to those who have many Pairs of *Canary-Birds*; and by means of this easy Method, you will never be dubious of the day your Little Ones are to come into the World; and you will pleasure yourself and your Friends, reading from time to time, the Number of Broods, and of Eggs you had of each Pair; how many have prov'd Addle, how many have Hatch'd, how many have Dy'd soon after; and in fine the exact Age of those you have from each Pair. Leave a white Leaf at certain Distances, to set down all the Remarks you make successively, which is the more pleasant, because it does not charge

charge the Memory, there being no more requir'd than the Booking down of those Particulars as they occur. This Method is absolutely necessary, in order to know the fix'd day on which the little Birds are to be Hatch'd, because the mistaking of a day sometimes occasions the loss of a whole Brood, by failing thereby to put into their Huts what is usual to be given them the day before they Hatch, whereof I have spoken at large in the 7th Chapter.

As for the time that *Canary-Birds* live, I cannot ascertain it, for as I have said, that the *Canary-Birds* differ in Constitution, it may be thence concluded, that they live longer or shorter, according to the Strength of their Constitution. But to be somewhat more particular; a Cock *Canary-Bird* that is put up to Breed every Year, seldom lives above 10 Years, and generally the 7th Year is not fit to be put up to Breed, for very often he will leave the Hen to Lay Addle Eggs.

A Hen that Sits every Year, seldom goes beyond six or seven Years, or if she does, it is because she has been well manag'd, or that she is, as has been

rest, so that they may not hear them Sing, otherwise it may happen, that your *Canary Bird* hearing the others that are in their Huts, may grow hot in love, and will Dye of that Distemper in a few days, for want of being Pair'd.

If you would proceed orderly, without burdening your Memory, which ought to be better employ'd than on the Thoughts of *Canary-Birds*, you must every Year make a little white Paper Book, to set down the precise day you gave the Hens their Eggs to Sit on; the day they are to Hatch, if good: I speak to those who have many Pairs of *Canary-Birds*; and by means of this easy Method, you will never be dubious of the day your Little Ones are to come into the World; and you will pleasure yourself and your Friends, reading from time to time, the Number of Broods, and of Eggs you had of each Pair; how many have prov'd Addle, how many have Hatch'd, how many have Dy'd soon after; and in fine the exact Age of those you have from each Pair. Leave a white Leaf at certain Distances, to set down all the Remarks you make successively, which is the more pleasant, because it does not charge

charge the Memory, there being no more requir'd than the Booking down of those Particulars as they occur. This Method is absolutely necessary, in order to know the fix'd day on which the little Birds are to be Hatch'd, because the mistaking of a day sometimes occasions the loss of a whole Brood, by failing thereby to put into their Huts what is usual to be given them the day before they Hatch, whereof I have spoken at large in the 7th Chapter.

As for the time that *Canary-Birds* live, I cannot ascertain it, for as I have said, that the *Canary-Birds* differ in Constitution, it may be thence concluded, that they live longer or shorter, according to the Strength of their Constitution. But to be somewhat more particular; a Cock *Canary-Bird* that is put up to Breed every Year, seldom lives above 10 Years, and generally the 7th Year is not fit to be put up to Breed, for very often he will leave the Hen to Lay Addle Eggs.

A Hen that Sits every Year, seldom goes beyond six or seven Years, or if she does, it is because she has been well manag'd, or that she is, as has been

been said, of an extraordinary strong Constitution.

Of all sorts of *Canary-Birds* the Grey live longest, being much stronger than the Copple Crown'd, Lemmon, &c.

A *Canary-Bird* of a good Constitution, and that has been well manag'd, that is, never put to Breed, may live 22 Years, but then he is subject to many Diseases, as Want of Stomach, Blindness, loosing his Claws and Voice, and sometimes to the Gout; and in that Condition his languishing Life, which threatens a speedy and certain Death, is as troublesom to him, by reason of the sharp Pains he continually endures, as a Mans is to him, when he is arriv'd at a certain uneasy degree of old Age, at which the only sovereign Remedy is no other than Death.

But enough of this Subject; tho' I have not yet exhausted it, yet I am apt to believe I have said enough to increase the Number of curious Persons, who have not ventur'd to give themselves the innocent and agreeable Diversi-
on of keeping *Canary-Birds*, for want of understanding them; as well in regard
to

to their good Qualities, as to their Faults and Difeates, and the feveral Accidents that befall them, during the Courfe of their Life, efpecially when they are defign'd to Breed.

In tenui Labor, at tenuis non Gloria.

FINIS.

BOOKS Printed for *Bernard Lintott.*

W <i>Iquefort's</i> Embassador, L. Pr.	1	15	0
———— Ditto Small Paper	1	5	0
Mr. <i>Popes</i> Homer, Vol. I. II. III. IV.	5	0	0
———— Ditto Small Paper	2	16	0
———— His Miscellaneous Works	1	5	0
———— Ditto, Small Paper	0	14	0
Mr. Serjeant <i>Lutwyche's</i> Reports, } Vol. I. II.	2	18	0
<i>Bobun</i> of the Rights of Boroughs	0	12	0
<i>Basneages</i> Hist. of the <i>Jews</i>	1	5	0
<i>Lawrence</i> of Gardening	0	1	0
———— Part II.	0	2	0
Mr. <i>Pointer's</i> Chronological Hist. 2 Vols.	0	5	0
Continuation the 1 st . Y. of King <i>George</i>	0	1	0
———— 2.	0	1	0
———— 3.	0	1	0
Poems by the Earl of <i>Rochester</i> and } Lord <i>Roscomon</i>	0	3	6
Poems on several Occasions by Mr. <i>Fenton</i>	0	3	6
Poems by his Grace of <i>Buckingham</i>	0	3	6
Poems entituled <i>Oxford</i> and <i>Camb.</i> Misc.	0	5	0
Poems in 4 Vols. 8 ^o .	0	18	0
Poems by the Lady <i>Chudleigh</i> 3 ^d Edit.	0	2	0
Poems by the great <i>Shakespear</i>	0	3	0
<i>Windfor Forest</i> 2 Edit.	0	1	0
<i>Essay on Criticism</i> 5 Edit.	0	1	0
<i>Rape of the Lock</i> 5 Edit.	0	1	0
By Mr. <i>Pope</i> & Temple of Fame 2 } Edit.	0	1	0
<i>Ode on St. Cecilia's</i> } Day	0	0	6
		By	

Books Printed for Bernard Lintot.

By Mr. Gay.	What d'ye call it, a Farce 2 Edit.	} 0	1	0
	Art of Walking the Streets 2 Edit.	} 0	1	0
	Three Hours after Marriage	} 0	1	6
	Wife of Bath	0	1	0
	Journey to Exeter	0	0	6
	Plays by the Rt. Honble Ld Lansdown	0	3	6
	Plays by Sr. Richard Steel	0	3	0
	Plays by Nich. Rowe, Esq; 2 Vol.	0	10	0
	Plays by Mr Southern 2 Vol.	0	5	0
By N. Rowe. Esq;	The Tragedy of Jane Shore 3 Edit.	} 0	1	6
	The Lady Jane Gray 2 Edit.	} 0	1	6
	Dennis's Remarks on Cato	0	1	0
	On Mr. Popes Essay on Criticism	0	0	6
	Essay on Publick Spirit	0	0	6
	On the Writings of Shakespear	0	1	0
	Grounds of Criticism	0	1	6
	Liberty Asserted, a Tragedy	0	1	6
	Appius and Virginia, a Tragedy	0	1	6
	Dr. Kings Miscellanies	0	5	0
	Art of Cookery 2 Edit.	0	1	6
	Art of Love	0	3	6
	Useful Transactions 3 Parts	0	3	0
	Useful Miscellanies	0	0	6
	La Motte's Remarks on Homer	0	1	0
	Roman Catholick System of Allegiance	0	1	0
	Religion of the Wits at Buttons	0	1	0
	Geddes Tract against Popery	0	5	0
	Homer by Mad. Dacier 5 Vol.	0	15	0
	Horace 2 Vol, Eng. and Lat.	0	10	0
	E. of Lauderdale's Virgil 2 Edit.	0	5	0
	Q. Curtius Englished by Mr. Digby	0	5	0
	C. Nepos Eng.	0	2	0
	Trapps Relectiones Poeticæ	0	3	0
	Drydens Art of Painting 2 Edit.	0	5	0
	Art of getting Pretty Children	0	1	6

Books Printed for Bernard Lintot.

<i>C. Gabala's</i> Diverting History	o	1	6
Mottos to the Tatler and Spectator	o	1	6
The invisible Mistress, by Mr. Cibber	o	1	o
Muscipula	o	o	6
<i>Moliere's</i> Comedys 6 Vols. Eng.	o	15	o
<i>Theophrastus's</i> Characters	o	4	o
Dr. King's Demy Gods 2 Edit.	o	2	o
<i>Du Pile</i> of mixing Colours	o	o	6
Cymbalum Mundi	o	1	o
Lives of the Popes	o	5	o
<i>Le Clercs</i> Abridgment of Clarendon	o	2	o
Hist. of <i>Formosa</i> 2 Edit.	o	5	o
— of <i>Holland</i> 2 Vols.	o	10	o
— of <i>Guinea</i>	o	6	o
— of <i>Saracens</i> , by Ockley 1 Vol.	o	5	o
— Ditto 2 Vol.	o	5	o
Capt Rogers Voyage to the South Sea 2 Edit.	} o	5	o
Capt. Cooks Voyage to Ditto 2 Vols.	o	10	o
<i>Perraults</i> Lives 2 Vols.	o	5	o
Mrs. <i>Phillips's</i> Letters	o	3	o
Dialogues of the Nature of the Gods	o	2	o
<i>Kennedy</i> of the Diseases of the Eyes	o	2	o
<i>Johnsons</i> Nouns and Verbs	o	1	o
— Decerpta ex Ovidij Fastis	o	1	o
— Ovids Metamorphosis	o	1	o
— Phedrus	o	1	o
Modern Policy	o	o	6
Monthly Catalogues of Books Publish'd	o	o	3
Managers Pro and Con	o	o	6
Annotations on the Tatlers	o	2	o
Conduct of the <i>Irish</i> Purse	o	o	6
Monthly Amusements, each	o	1	o
Votes of Parliament 1714, 15, 16, 17			
Report of the Secret Committee	o	3	6
General <i>Stanhopes</i> Answer to the Re- port of the Commissioners.	} o	o	6
<i>Bohuns</i> Institutio Legalis 2 Edit.	o	6	o
Complete Court Keeper 2 Edit.	o	6	o

Court

Books Printed for *Bernard Lintot.*

Court Keepers Companion	o	2	6
Accomplish'd Conveyancer 3 Vols.	o	18	o
Clerks Remembrancer	o	3	6
Grand Precedent for Conveyancing	o	6	o
Modern Justice 2 Edit.	o	5	o
Appendix to the Modern Justice	o	1	6
<i>Minwoods</i> Forest Laws	o	5	o
<i>Nelsons</i> Reports in Chancery	o	3	6
<i>Noy's</i> Rights o'th' Crown	o	2	o
Ld. Chief Justice <i>Egerton</i> against <i>Cooks</i> Reports	} o	1	6
A Table to the Statutes	o	o	6
Discourse on Treason and Bills of Attainder	} o	1	6
Journals of the House of Commons	o	3	o
Rules of Government, by Sr. <i>John Heywood</i>	} o	1	o
Diseases incident to Tradesmen	o	3	6
Advice to young Gentlemen	o	1	6
<i>Du Pins</i> Abbrev. of Church Hist. 4 Vols.	o	10	o
Bp. <i>Williams's</i> Sermons, <i>Boyles</i> Lecture	o	5	o
Dr. <i>Tillys</i> Sermons	o	5	o
—— his Devotions	o	2	o
Mr. <i>Fidde's</i> Body of Divinity Vol. 1.	1	5	o
—— on the Sacrament	o	1	o
Dean <i>Gardiner</i> on the Beatitudes	o	6	o
—— Translation of <i>Rapin de Hortis</i>	o	3	6
Dr. <i>Goddards</i> Six Sermons	o	1	o
Mr. <i>Lambs</i> Abridgment of <i>Bishop Pearson</i> on the Creed	} o	2	6
<i>Timothy</i> and <i>Philatheus</i> 3 Vols.	o	15	o
Essay on Private Judgment	o	3	o
<i>Seneca's</i> Morals 11th Edit.	o	5	o
<i>Plato's</i> Immortality of the Soul	o	1	o
<i>Thorp's</i> Assize Sermon	o	o	4
<i>Laurence's</i> Assize Sermon	o	o	4
Sermon at the Funeral of a Day Labourer	o	o	4
<i>Bryar's</i> Catechism	o	o	6
—— Prayers and Meditations	o	o	3
<i>Trapp</i> on the Lord's Prayer	o	1	o

Peck

Books Printed for Bernard Lintot.

<i>Peck on the Creation</i>	o	I	o
<i>Setvens's Sermon against Popery</i>	o	o	4
<i>Mesech and Kedar, v. Trapp</i>	o	o	4
<i>Jacksons Assize Sermon</i>	o	o	4
<i>Dr. Goddard's Sermon on Censure</i>	o	o	4
— on Gods Mercies	o	o	4
<i>Lushington's Sermon</i>	o	o	4
<i>Dibbons's Sermon in Holland</i>	o	o	4
<i>Dubourdieu's Sermon</i>	o	o	4
<i>Chillingworths Judgment</i>	o	o	4
<i>Divinity and Law reconcil'd</i>	o	I	o
<i>State of Trinity College in Cambridge</i>	o	o	4
<i>Sentences of Ali by the Rev. Mr Ockley</i>	o	o	6
<i>Lambs Sermons, each</i>	o	o	4
<i>Toland's Letter against Popery</i>	o	o	6
— Letters to <i>Serena</i>	o	o	6
— <i>Anglia Libera</i>	o	3	6
— <i>Art of Governing by Partys</i>	o	3	o
— <i>Vindicius Liberus</i>	o	2	6
— <i>Art of Restoring</i>	o	o	6
— <i>General Monks Letter</i>	o	I	o
— <i>Reasons for Naturalizing the Jews</i>	} o	I	o
— <i>Grand Mystery laid open</i>	o	o	6
— <i>Amyntor</i>	o	2	6
<i>A Treatise on the Small Pox</i>	o	I	o
<i>Recruiting Officer, a Comedy 5 Edit.</i>	o	I	o
<i>Beaux Stratagem, a Comedy 5 Edit.</i>	o	I	o
<i>Busy Body 2 Edit.</i>	o	I	o
<i>Quaker of Deale 2 Edit.</i>	o	I	o
<i>Perkin Warbeck 2 Edit.</i>	o	I	o
<i>Phædra and Hippolitus</i>	o	I	6
By <i>Venus & Adonis, a Masque</i>	o	o	6
Mr. Cibber. <i>Myrtillo, a Masque</i>	o	o	6
<i>Loves Last Shift</i>	o	I	o
<i>Ladys Last Stake</i>	o	I	6
<i>Double Gallant</i>	o	I	6
<i>Perrolla and Izzadora</i>	o	I	6
<i>Comical Lover</i>	o	I	6
<i>The Non-juror</i>	o	I	6